

A Primal-Dual SGD Algorithm for Distributed Nonconvex Optimization

Xinlei Yi, Shengjun Zhang, Tao Yang, Tianyou Chai, *Fellow, IEEE*, and Karl Henrik Johansson, *Fellow, IEEE*

Abstract—The distributed nonconvex optimization problem of minimizing a global cost function formed by a sum of n local cost functions by using local information exchange is considered. This problem is an important component of many machine learning techniques with data parallelism, such as deep learning and federated learning. We propose a distributed primal-dual stochastic gradient descent (SGD) algorithm, suitable for arbitrarily connected communication networks and any smooth (possibly nonconvex) cost functions. We show that the proposed algorithm achieves the linear speedup convergence rate $O(1/\sqrt{nT})$ for general nonconvex cost functions and the linear speedup convergence rate $O(1/(nT))$ when the global cost function satisfies the Polyak-Lojasiewicz (P-L) condition, where T is the total number of iterations. We also show that the output of the proposed algorithm with constant parameters linearly converges to a neighborhood of a global optimum. We demonstrate through numerical experiments the efficiency of our algorithm in comparison with the baseline centralized SGD and recently proposed distributed SGD algorithms.

Index Terms—Distributed nonconvex optimization, linear speedup, Polyak-Lojasiewicz (P-L) condition, primal-dual algorithm, stochastic gradient descent.

I. INTRODUCTION

CONSIDER a network of n agents, each of which has a local smooth (possibly nonconvex) cost function $f_i: \mathbb{R}^p \rightarrow \mathbb{R}$. All agents collaboratively solve the following optimization problem:

$$\min_{x \in \mathbb{R}^p} f(x) := \frac{1}{n} \sum_{i=1}^n f_i(x). \quad (1)$$

Each agent i only has information about its local cost function f_i and communicates with its neighbors through the

Manuscript received February 11, 2022; accepted February 25, 2022. This work was supported by the Knut and Alice Wallenberg Foundation, the Swedish Foundation for Strategic Research, the Swedish Research Council, and the National Natural Science Foundation of China (62133003, 61991403, 61991404, 61991400). Recommended by Associate Editor Qing-Long Han. (Corresponding author: Tao Yang.)

Citation: X. L. Yi, S. J. Zhang, T. Yang, T. Y. Chai, and K. H. Johansson, "A primal-dual SGD algorithm for distributed nonconvex optimization," *IEEE/CAA J. Autom. Sinica*, vol. 9, no. 5, pp. 812–833, May 2022.

X. L. Yi and K. H. Johansson are with the Division of Decision and Control Systems, School of Electrical Engineering and Computer Science, KTH Royal Institute of Technology, and also affiliated with the Digital Futures, Stockholm 10044, Sweden (e-mail: xinleiy@kth.se; kallej@kth.se).

S. J. Zhang is with the Department of Electrical Engineering, University of North Texas, Denton, TX 76203 USA (e-mail: ShengjunZhang@my.unt.edu).

T. Yang and T. Y. Chai are with the State Key Laboratory of Synthetical Automation for Process Industries, Northeastern University, Shenyang 110819, China (e-mail: yangtao@mail.neu.edu.cn; tychai@mail.neu.edu.cn).

Color versions of one or more of the figures in this paper are available online at <http://ieeexplore.ieee.org>.

Digital Object Identifier 10.1109/JAS.2022.105554

underlying communication network. The communication network is modeled by an undirected graph $\mathcal{G} = (\mathcal{V}, \mathcal{E})$, where $\mathcal{V} = \{1, \dots, n\}$ is the agent set, $\mathcal{E} \subseteq \mathcal{V} \times \mathcal{V}$ is the edge set, and $(i, j) \in \mathcal{E}$ if agents i and j communicate with each other. The set $\mathcal{N}_i = \{j \in \mathcal{V} : (i, j) \in \mathcal{E}\}$ is the neighboring set of agent i . The optimization problem (1) incorporates many popular machine learning approaches with data parallelism, such as deep learning [1] and federated learning [2]. A star graph is a special undirected graph, in which there is one and only one agent (hub agent) that connects to all of the other agents (leaf agents) and each leaf agent only connects to the hub agent. Such a graph corresponds to the master/worker architecture adopted by many parallel learning algorithms.

Due to the nonconvexity, convergence results typically ensure that the distributed algorithms find first-order stationary points

$$\{x \in \mathbb{R}^p : \nabla f(x) = \mathbf{0}_p\}$$

which could be local maxima or minima. Global optima are usually hard to find. For example, in [3]–[9], it was shown that first-order stationary points can be found with an $O(1/T)$ convergence rate, i.e.,

$$\frac{1}{T} \sum_{k=0}^{T-1} \|\nabla f(\bar{x}_k)\|^2 = O\left(\frac{1}{T}\right)$$

where T is the total number of iterations, ∇f denotes the gradient of f , and $\bar{x}_k = \frac{1}{n} \sum_{i=1}^n x_{i,k}$ with $x_{i,k} \in \mathbb{R}^p$ being agent i 's estimate of the optimal solution at time instant k .

Note that the algorithms proposed in the aforementioned references use at least gradient information of the cost functions, and sometimes even the second- or higher-order information. However, in many applications explicit expressions of the gradients are often unavailable or difficult to obtain. In this paper, we consider the case where each agent can only collect stochastic gradients of its local cost function and propose a distributed stochastic gradient descent (SGD) algorithm to solve (1). In general, SGD algorithms are suitable for scenarios where explicit expressions of the gradients are unavailable or difficult to obtain. For example, in some big data applications, such as empirical risk minimization, the actual gradient is calculated from the entire data set, which results in a heavy computational burden. A stochastic gradient can be calculated from a randomly selected subset of the data and is often an efficient way to replace the actual gradient. Other examples which SGD algorithms are suitable for include scenarios where data are arriving sequentially such as in online learning [10].

A. Literature Review

When the communication network is a star graph, various parallel SGD algorithms have been proposed to solve (1). A potential performance bottleneck of such algorithms lies on the communication burden of the master. To overcome this issue, a promising strand of research is combining parallel SGD algorithms with communication reduction approaches, e.g., asynchronous parallel SGD algorithms [11]–[15], gradient compression based parallel SGD algorithms [12], [16]–[19], periodic averaging based parallel SGD algorithms [17], [18], [20]–[24], and parallel SGD algorithm with dynamic batch sizes [25]. Convergence properties of these algorithms have been analyzed in detail. In particular, in [17], [21], [23], [25], an $O(1/\sqrt{nT})$ convergence rate has been established for general nonconvex cost functions. This rate is n times faster than the well known $O(1/\sqrt{T})$ convergence rate established by SGD over a single agent, and thus a linear speedup in the number of agents is achieved. In [24], [25], the convergence rate has been improved to $O(1/(nT))$ when the global cost function satisfies the P-L condition, which also achieves a linear speedup. In addition to the star architecture restriction, aforementioned parallel SGD algorithms require certain restrictions on the cost functions, such as bounded gradients of the local cost functions or bounded difference between the gradients of the local and global cost functions.

Distributed algorithms executed over arbitrarily connected communication networks have been suggested to overcome communication bottlenecks for parallel SGD algorithms. Various distributed SGD algorithms have been proposed to solve (1), e.g., synchronous distributed SGD algorithms [23], [26]–[28], asynchronous distributed SGD algorithms [29], [30], compression based distributed SGD algorithms [31]–[34], and periodic averaging based distributed SGD algorithm [35]. Convergence properties of these algorithms have been analyzed and the linear speedup convergence rate $O(1/\sqrt{nT})$ has been established for general nonconvex cost functions [23], [27], [30], [31], [33]–[35]. However, similarly to aforementioned parallel SGD algorithms, these distributed algorithms require restrictive assumptions on the cost functions. In order to remove these restrictions, the authors of [36] proposed a variant of the distributed SGD algorithm proposed in [27], named D^2 , in which each agent stores the stochastic gradient and its local model in last iteration and linearly combines them with the current stochastic gradient and local model. For this algorithm the authors established the linear speedup convergence rate $O(1/\sqrt{nT})$, but they required that the eigenvalues of the mixing matrix associated with the communication network are strictly greater than $-1/3$. The authors of [37], [38] proposed distributed stochastic gradient tracking algorithms suitable for arbitrarily connected communication networks. However, these algorithms only achieve $O(1/\sqrt{T})$ convergence rate, which is not a speedup. Moreover, gradient tracking algorithms have the common potential drawback that in order to track the global gradient, at each iteration each agent needs to communicate one additional p -dimensional variable with its neighbors. This results in heavy communication burden when p is large. Note that all

aforementioned distributed SGD algorithms converge to stationary points, which may be local or global optima, or saddle points. None of existing studies on distributed SGD algorithms consider finding the global optimum when the global cost function satisfies some additional property, such as the P-L condition studied for the parallel algorithms in [24], [25]. Noting above, two core theoretical questions arise.

Q1) Are there any distributed SGD algorithms that are not only suitable for arbitrarily connected communication networks and any smooth cost functions but also find stationary points with the linear speedup convergence rate $O(1/\sqrt{nT})$?

Q2) If the P-L condition holds in addition, can the above SGD algorithms find the global optimum with the linear speedup convergence rate $O(1/(nT))$ as achieved in [24], [25]?

B. Main Contributions

This paper provides positive answers for the above two questions. More specifically, the contributions of this paper are summarized as follows.

i) We propose a distributed primal-dual SGD algorithm to solve the optimization problem (1). In the proposed algorithm, each agent maintains the primal and dual variable sequences and only communicates the primal variable with its neighbors. This algorithm is suitable for arbitrarily connected communication networks and any smooth (possibly nonconvex) cost functions.

ii) We show in Corollary 1 that our algorithm finds a stationary point with the linear speedup convergence rate $O(1/\sqrt{nT})$ for general nonconvex cost functions, thus answers Q1). Compared with [17], [21], [23], [25], [27], [30], [31], [33]–[36], we achieve the same convergence rate but under weaker assumptions related to network architectures and/or cost functions, and compared with [37], [38], we not only establish linear speedup but also just use half communication in each iteration.

iii) We show in Theorem 3 that our algorithm finds a global optimum with the linear speedup convergence rate $O(1/(nT))$ when the global cost function satisfies the P-L condition, thus answers Q2). Compared with [24], [25], [34], [39]–[41], we achieve the same convergence rate but under weaker assumptions related to network architectures and/or cost functions, and compared with [18], [26], [42]–[46], we not only establish linear speedup but also relax the strong convexity by the P-L condition.

iv) We show in Theorems 4 and 5 that the output of our algorithm with constant parameters linearly converges to a neighborhood of a global optimum when the global cost function satisfies the P-L condition. Compared with [26], [46]–[49], which used the strong convexity assumption, we achieve the similar convergence result under weaker assumptions on the cost function.

The detailed comparison of this paper with other related studies in the literature is summarized in Table I.

C. Outline

The rest of this paper is organized as follows. Section II presents the novel distributed primal-dual SGD algorithm.

TABLE I
COMPARISON OF THIS PAPER TO SOME RELATED WORKS

Reference	Problem type	Extra assumption	Communication network	Communicated variable	Communication rounds	Convergence rate
[17]	Nonconvex	Bounded $\ \nabla f_i - \nabla f\ $	Star graph	One quantized variable	$O(n^{5/4}T^{3/4})$	$O(1/\sqrt{nT})$
[18]	Nonconvex	Identical ∇f_i	Star graph	One quantized variable	$O(T)$	$O(1/\sqrt{T})$
	Strongly convex					$O(1/T)$
[21]	Nonconvex	Bounded $\ \nabla f_i\ $	Star graph	One full-information variable	$O(n^{3/4}T^{3/4})$	$O(1/\sqrt{nT})$
[23]	Nonconvex	Bounded $\ \nabla f_i - \nabla f\ $	Star graph	Two full-information variables	$O(n^{3/4}T^{3/4})$	$O(1/\sqrt{nT})$
			Connected graph		$O(T)$	
[24]	P-L condition	Identical ∇f_i	Star graph	One full-information variable	$O((nT)^{1/3})$	$O(1/(nT))$
[25]	Nonconvex	Identical ∇f_i , exponentially increasing batch size	Star graph	One full-information variable	$O(\sqrt{nT} \log(\frac{T}{n}))$	$O(1/\sqrt{nT})$
	P-L condition				$O(\log(T))$	$O(1/(nT))$
[26]	Nonconvex	Bounded $\ \nabla f_i\ $	Connected graph	One full-information variable	$O(T)$	$O(1/T^\theta), \forall \theta \in (0, 0.5)$
	Strongly convex					$O(1/T)$; linearly to a neighbor of the global optimum (constant stepsize)
[27]	Nonconvex	Bounded $\ \nabla f_i - \nabla f\ $	Connected graph	One full-information variable	$O(T)$	$O(1/\sqrt{nT})$
[30]	Nonconvex	Bounded $\ \nabla f_i - \nabla f\ $	Uniformly jointly strongly connected digraph	One full-information variable	$O(T)$	$O(1/\sqrt{nT})$
[31]	Nonconvex	Bounded $\ \nabla f_i - \nabla f\ $	Connected graph	One compressed variable	$O(T)$	$O(1/\sqrt{nT})$
[33]	Nonconvex	Bounded $\ \nabla f_i\ $	Strongly connected digraph	One quantized variable	$O(T)$	$O(1/\sqrt{nT})$
[34]	Nonconvex	Bounded $\ \nabla f_i\ $	Connected graph	One compressed variable	Event-triggered	$O(1/\sqrt{nT})$
	Strongly convex					$O(1/(nT))$
[35]	Nonconvex	Identical ∇f_i	Connected graph	One full-information variable	$O(n^{3/2}\sqrt{T})$	$O(1/\sqrt{nT})$
[36]	Nonconvex	The eigenvalues of the mixing matrix are strictly greater than $-1/3$	Connected graph	One full-information variable	$O(T)$	$O(1/\sqrt{nT})$
[37], [38]	Nonconvex	No	Connected graph	Two full-information variables	$O(T)$	$O(1/\sqrt{T})$
[39]	Strongly convex	Bounded $\ \nabla f_i\ $	Star graph	One full-information variable	$O(\sqrt{T/n})$	$O(1/(nT))$
[40]	Strongly convex	Bounded $\ \nabla f_i\ $	Connected graph	One compressed variable	$O(T)$	$O(1/(nT))$
[41]	Strongly convex	No	Connected graph	Two full-information variables	$O(T)$	$O(1/(nT))$
[42]	Strongly convex	Identical ∇f_i	Connected graph	One full-information variable	$O(T)$	$O(1/T)$
[43]	Strongly convex	No	Connected graph	One full-information variable	$O(\sqrt{T})$	$O(1/T)$
[44]	Strongly convex	Bounded $\ \nabla f_i\ $	Uniformly jointly strongly connected digraph	One full-information variable	$O(T)$	$O(1/T)$
[45]	Strongly convex	No	Connected graph in expectation	One full-information variable	$O(T)$	$O(1/T)$
[46]	Strongly convex	No	Connected graph	One full-information variable	$O(T)$	$O(1/T)$; linearly to a neighbor of the global optimum (constant stepsize)
[47]	Strongly convex	No	Connected graph	One full-information variable	$O(T)$	Linearly to a neighbor of the global optimum (constant stepsize)
[48]	Strongly convex	No	Connected graph	Two full-information variables	$O(T)$	Linearly to a neighbor of the global optimum (constant stepsize)
[49]	Strongly convex	No	Strongly connected digraph	Two full-information variables	$O(T)$	Linearly to a neighbor of the global optimum (constant stepsize)
This paper	Nonconvex	No	Connected graph	One full-information variable	$O(T)$	$O(1/\sqrt{nT})$
	P-L condition					$O(1/(T^\theta)), \forall \theta \in (0, 1)$; linearly to a neighbor of the global optimum (constant stepsize)
	Bounded f_i^*					$O(1/(nT))$

Section III analyzes its convergence rate. Numerical experiments are given in Section IV. Finally, concluding remarks are offered in Section V. To improve the readability, all the proofs are given in the appendix.

Notations: \mathbb{N}_0 and \mathbb{N}_+ denote the set of nonnegative and positive integers, respectively. $[n]$ denotes the sets $\{1, \dots, n\}$ for any $n \in \mathbb{N}_+$. $\mathbf{1}_n$ ($\mathbf{0}_n$) denotes the column one (zero) vector of dimension n . $\text{col}(z_1, \dots, z_k)$ is the concatenated column vector of vectors $z_i \in \mathbb{R}^{p_i}$, $i \in [k]$. $\|\cdot\|$ represents the Euclidean norm for vectors or the induced 2-norm for matrices.

II. DISTRIBUTED PRIMAL-DUAL SGD ALGORITHM

In this section, we propose a novel distributed SGD algorithm based on the primal-dual method.

Denote $\mathbf{x} = \text{col}(x_1, \dots, x_n)$, $\tilde{f}(\mathbf{x}) = \sum_{i=1}^n f_i(x_i)$, and $\mathbf{L} = L \otimes \mathbf{I}_p$, where $L = (L_{ij})$ is the weighted Laplacian matrix associated with the undirected communication graph \mathcal{G} . Recall that the Laplacian matrix L is positive semi-definite and $\text{null}(L) = \{\mathbf{1}_n\}$ when \mathcal{G} is connected [50]. The optimization problem (1) is equivalent to the following constrained optimization problem:

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^{np}} \quad & \tilde{f}(\mathbf{x}) \\ \text{s.t.} \quad & \mathbf{L}^{1/2} \mathbf{x} = \mathbf{0}_{np}. \end{aligned} \quad (2)$$

Here, $\mathbf{L}^{1/2} = L^{1/2} \otimes \mathbf{I}_p$ and $L^{1/2}$ is the square root of the positive semi-definite matrix L . Moreover, $\mathbf{L}^{1/2} \mathbf{x} = \mathbf{0}_{np}$ is equivalent to $\mathbf{x} = \mathbf{1}_n \otimes \mathbf{x}$ due to the fact that $\text{null}(L^{1/2}) = \text{null}(L) = \{\mathbf{1}_n\}$.

Let $\mathbf{u} \in \mathbb{R}^{np}$ denote the dual variable. Then the augmented Lagrangian function associated with (2) is

$$\mathcal{A}(\mathbf{x}, \mathbf{u}) = \tilde{f}(\mathbf{x}) + \frac{\alpha}{2} \mathbf{x}^T \mathbf{L} \mathbf{x} + \beta \mathbf{u}^T \mathbf{L}^{1/2} \mathbf{x} \quad (3)$$

where $\alpha > 0$ and $\beta > 0$ are parameters to be designed later.

Based on the primal-dual gradient method, a distributed SGD algorithm to solve (2) is

$$\mathbf{x}_{k+1} = \mathbf{x}_k - \eta_k (\alpha_k \mathbf{L} \mathbf{x}_k + \beta_k \mathbf{L}^{1/2} \mathbf{u}_k + \mathbf{g}_k^u) \quad (4a)$$

$$\mathbf{u}_{k+1} = \mathbf{u}_k + \eta_k \beta_k \mathbf{L}^{1/2} \mathbf{x}_k, \quad \forall \mathbf{x}_0, \mathbf{u}_0 \in \mathbb{R}^{np} \quad (4b)$$

where $\eta_k > 0$ is the stepsize at iteration k , $\alpha_k > 0$ and $\beta_k > 0$ are the values of the parameters α and β at iteration k , respectively, and $\mathbf{g}_k^u = \text{col}(g_{1,k}^u, \dots, g_{n,k}^u)$ with $g_{i,k}^u = g_i(x_{i,k}, \xi_{i,k})$ being the stochastic gradient of f_i at $x_{i,k}$ and $\xi_{i,k}$ being a random variable. Denote $\mathbf{v}_k = \text{col}(v_{1,k}, \dots, v_{n,k}) = \mathbf{L}^{1/2} \mathbf{u}_k$. Then the recursion (4) can be rewritten as

$$\mathbf{x}_{k+1} = \mathbf{x}_k - \eta_k (\alpha_k \mathbf{L} \mathbf{x}_k + \beta_k \mathbf{v}_k + \mathbf{g}_k^u) \quad (5a)$$

$$\mathbf{v}_{k+1} = \mathbf{v}_k + \eta_k \beta_k \mathbf{L} \mathbf{x}_k, \quad \forall \mathbf{x}_0 \in \mathbb{R}^{np}, \sum_{j=1}^n v_{j,0} = \mathbf{0}_p. \quad (5b)$$

The initialization condition $\sum_{j=1}^n v_{j,0} = \mathbf{0}_p$ is derived from $\mathbf{v}_0 = \mathbf{L}^{1/2} \mathbf{u}_0$, and it is easy to be satisfied, for example, $v_{i,0} = \mathbf{0}_p$, $\forall i \in [n]$, or $v_{i,0} = \sum_{j=1}^n L_{ij} x_{j,0}$, $\forall i \in [n]$. Note that (5) can be written agent-wise as

$$x_{i,k+1} = x_{i,k} - \eta_k \left(\alpha_k \sum_{j=1}^n L_{ij} x_{j,k} + \beta_k v_{i,k} + g_{i,k}^u \right) \quad (6a)$$

$$v_{i,k+1} = v_{i,k} + \eta_k \beta_k \sum_{j=1}^n L_{ij} x_{j,k},$$

$$\forall x_{i,0} \in \mathbb{R}^p, v_{i,0} = \mathbf{0}_p, \forall i \in [n]. \quad (6b)$$

This corresponds to our proposed distributed primal-dual SGD algorithm, which is presented in pseudo-code as Algorithm 1.

Algorithm 1 Distributed Primal-Dual SGD Algorithm

- 1: **Input:** parameters $\{\alpha_k\}, \{\beta_k\}, \{\eta_k\} \subseteq (0, +\infty)$.
 - 2: **Initialize:** $x_{i,0} \in \mathbb{R}^p$ and $v_{i,0} = \mathbf{0}_p$, $\forall i \in [n]$.
 - 3: **for** $k = 0, 1, \dots$ **do**
 - 4: **for** $i = 1, \dots, n$ in parallel **do**
 - 5: Broadcast $x_{i,k}$ to \mathcal{N}_i and receive $x_{j,k}$ from $j \in \mathcal{N}_i$;
 - 6: Sample stochastic gradient $g_i(x_{i,k}, \xi_{i,k})$;
 - 7: Update $x_{i,k+1}$ by (6a);
 - 8: Update $v_{i,k+1}$ by (6b).
 - 9: **end for**
 - 10: **end for**
 - 11: **Output:** $\{x_k\}$.
-

It should be pointed out that $\{\alpha_k\}, \{\beta_k\}, \{\eta_k\}, \mathbf{x}_0, \mathbf{v}_0$, and \mathbf{v}_1 in Algorithm 1 are deterministic, while $\{x_k\}_{k \geq 1}$ and $\{v_k\}_{k \geq 2}$ are random variables generated by Algorithm 1. Let \mathfrak{F}_k denote the σ -algebra generated by the random variables $\xi_{1,k}, \dots, \xi_{n,k}$ and let $\mathcal{F}_k = \bigcup_{s=1}^k \mathfrak{F}_s$. It is straightforward to see that x_k and v_{k+1} depend on \mathcal{F}_{k-1} and are independent of \mathfrak{F}_s for all $s \geq k$.

III. CONVERGENCE RATE ANALYSIS

In this section, we analyze the convergence rate of Algorithm 1. The following assumptions are made.

Assumption 1: The undirected communication graph \mathcal{G} is connected.

Assumption 2: The minimum function value of the optimization problem (1) is finite.

Assumption 3: Each local cost function f_i is smooth with constant $L_f > 0$, i.e.,

$$\|\nabla f_i(x) - \nabla f_i(y)\| \leq L_f \|x - y\|, \quad \forall x, y \in \mathbb{R}^p. \quad (7)$$

Assumption 4: The random variables $\{\xi_{i,k}, i \in [n], k \in \mathbb{N}_0\}$ are independent of each other.

Assumption 5: The stochastic estimate $g_i(x, \xi_{i,k})$ is unbiased, i.e., for all $i \in [n]$, $k \in \mathbb{N}_0$, and $x \in \mathbb{R}^p$,

$$\mathbf{E}_{\xi_{i,k}} [g_i(x, \xi_{i,k})] = \nabla f_i(x). \quad (8)$$

Assumption 6: The stochastic estimate $g_i(x, \xi_{i,k})$ has bounded variance, i.e., there exists a constant σ such that for all $i \in [n]$, $k \in \mathbb{N}_0$, and $x \in \mathbb{R}^p$,

$$\mathbf{E}_{\xi_{i,k}} [\|g_i(x, \xi_{i,k}) - \nabla f_i(x)\|^2] \leq \sigma^2. \quad (9)$$

Remark 1: The bounded variance assumption (Assumption 6) is weaker than the bounded second moment (or bounded gradient) assumption made in [11]–[13], [15], [19], [21], [26], [28], [33], [34], [39], [40], [44], [51]. Moreover, note that we make no assumption on the boundedness of the deviation between the gradients of local cost functions. In other words, we do not assume that $\frac{1}{n} \sum_{i=1}^n \|\nabla f_i(x) - \nabla f(x)\|^2$ is uniformly bounded, a common assumption made in studies of deep

learning, e.g., [17], [21], [23], [27], [29]–[31]. Also, we do not assume that the mean of each local stochastic gradient is the gradient of the global cost function, i.e., $\mathbf{E}_{\xi}[g_i(x, \xi)] = \nabla f(x)$, $\forall x \in \mathbb{R}^p$, $\forall i \in [n]$, which is commonly assumed in studies of empirical risk minimization and stochastic optimization, e.g., [14], [16], [18], [20], [22], [24], [25], [32], [35], [42].

A. Find Stationary Points

Let us consider the case when Algorithm 1 is able to find stationary points. We have the following convergence results.

Theorem 1: Suppose Assumptions 1–6 hold. Let $\{x_k\}$ be the sequence generated by Algorithm 1 with

$$\alpha_k = \kappa_1 \beta_k, \beta_k = \beta, \eta_k = \frac{\kappa_2}{\beta_k}, \forall k \in \mathbb{N}_0 \quad (10)$$

where $\kappa_1 > c_1$, $\kappa_2 \in (0, c_2(\kappa_1))$, and $\beta \geq c_0(\kappa_1, \kappa_2)$ with $c_0(\kappa_1, \kappa_2)$, c_1 , $c_2(\kappa_1) > 0$ defined in Appendix B. Then, for any $T \in \mathbb{N}_+$,

$$\frac{1}{T} \sum_{k=0}^{T-1} \mathbf{E} \left[\frac{1}{n} \sum_{i=1}^n \|x_{i,k} - \bar{x}_k\|^2 \right] = \mathcal{O}\left(\frac{1}{T}\right) + \mathcal{O}\left(\frac{1}{\beta^2}\right) \quad (11a)$$

$$\begin{aligned} \frac{1}{T} \sum_{k=0}^{T-1} \mathbf{E} [\|\nabla f(\bar{x}_k)\|^2] &= \mathcal{O}\left(\frac{\kappa_2 \beta}{T}\right) + \mathcal{O}\left(\frac{4\kappa_2}{n\beta}\right) \\ &+ \mathcal{O}\left(\frac{1}{T}\right) + \mathcal{O}\left(\frac{1}{\beta^2}\right) \end{aligned} \quad (11b)$$

$$\mathbf{E}[f(\bar{x}_T)] - f^* = \mathcal{O}(1) + \mathcal{O}\left(\frac{T}{n\beta^2}\right) + \mathcal{O}\left(\frac{T}{\beta^3}\right) \quad (11c)$$

where f^* is the minimum function value of the optimization problem (1) and $\bar{x}_k = \frac{1}{n} \sum_{i=1}^n x_{i,k}$.

Proof: The explicit expressions of the right-hand sides of (11a)–(11c) and the proof are given in Appendix B. It should be highlighted that the omitted constants in the first two terms on the right-hand side of (11b) do not depend on any parameters related to the communication network. ■

Noting the right-hand side of (11b) and $f(\bar{x}_0) - f^* = \mathcal{O}(1)$, the linear speedup (w.r.t. number of agents) can be established if we set $\beta = \kappa_2 \sqrt{T} / \sqrt{n}$, as shown in the following.

Corollary 1 (Linear Speedup): Under the same assumptions as in Theorem 1, let $\beta = \kappa_2 \sqrt{T} / \sqrt{n}$. Then, for any $T > \max\{n(c_0(\kappa_1, \kappa_2)/\kappa_2)^2, n^3\}$,

$$\frac{1}{T} \sum_{k=0}^{T-1} \mathbf{E} \left[\frac{1}{n} \sum_{i=1}^n \|x_{i,k} - \bar{x}_k\|^2 \right] = \mathcal{O}\left(\frac{n}{T}\right) \quad (12a)$$

$$\frac{1}{T} \sum_{k=0}^{T-1} \mathbf{E} [\|\nabla f(\bar{x}_k)\|^2] = \mathcal{O}\left(\frac{1}{\sqrt{nT}}\right) + \mathcal{O}\left(\frac{n}{T}\right) \quad (12b)$$

$$\mathbf{E}[f(\bar{x}_T)] - f^* = \mathcal{O}(1). \quad (12c)$$

Remark 2: It should be highlighted that the omitted constants on the first term in the right-hand side of (12b) do not depend on any parameters related to the communication network. The same linear speedup result as in (12b) was also established by the SGD algorithms proposed in [17], [21], [23], [25], [27], [30], [31], [33]–[36]. However, in [17], [23], [27], [30], [31], the additional assumption that the deviation

between the gradients of local cost functions is bounded was made; in [21], [33], [34], it was required that each local stochastic gradient has bounded second moment; in [25], [35], it was assumed that the mean of each local stochastic gradient is the gradient of the global cost function; and in [36], it was required that the eigenvalues of the mixing matrix are strictly greater than $-1/3$. Moreover, the algorithms proposed in [17], [25] are restricted to a star graph; the distributed momentum SGD algorithm proposed in [23] requires each agent i to communicate one additional p -dimensional variable besides the communication of $x_{i,k}$ with its neighbors at each iteration; and the algorithm proposed in [25] requires an exponentially increasing batch size, which is not favorable in practice. Under the same conditions, the well known $\mathcal{O}(1/\sqrt{T})$ convergence rate, which is not a speedup, was achieved by the distributed stochastic gradient tracking algorithm proposed in [37], [38]. Moreover, similarly to the distributed momentum SGD algorithm proposed in [23], one potential drawback of the distributed stochastic gradient tracking algorithms is that at each iteration each agent needs to communicate one additional variable. The potential drawbacks of the results stated in Corollary 1 are that i) we do not consider communication efficiency, which was considered in [17], [21], [25], [31], [33]–[35]; and ii) we consider undirected graphs rather than directed graphs as considered in [30], [33]. We leave the extension to the (time-varying) directed graphs with communication efficiency as future research directions.

B. Find Global Optimum

Let us next consider the case when Algorithm 1 finds global optimum. The following assumption is crucial.

Assumption 7: The global cost function $f(x)$ satisfies the Polyak-Łojasiewicz (P-Ł) condition with constant $\nu > 0$, i.e.,

$$\frac{1}{2} \|\nabla f(x)\|^2 \geq \nu(f(x) - f^*), \forall x \in \mathbb{R}^p. \quad (13)$$

It is straightforward to see that every (essentially or weakly) strongly convex function satisfies the P-Ł condition. The P-Ł condition implies that every stationary point is a global minimizer. But unlike (essentially or weak) strong convexity, the P-Ł condition alone does not imply convexity of f . Moreover, it does not imply that the global minimizer is unique either [52], [53].

Various practical applications, such as least squares and logistic regression, do not always have strongly convex cost functions. The cost function in least squares problems has the form

$$f(x) = \frac{1}{2} \|Ax - b\|^2$$

where $A \in \mathbb{R}^{m \times p}$ and $b \in \mathbb{R}^m$. Note that if A has full column rank, then $f(x)$ is strongly convex. However, if A is rank deficient, then $f(x)$ is not strongly convex, but it is convex and satisfies the P-Ł condition. The function $f(x) = x^2 + 3 \sin^2(x)$ given in [52] is an example of a nonconvex function satisfying the P-Ł condition with $\nu = 1/32$. Moreover, it was shown in [54] that the loss functions in some applications satisfy the P-Ł condition in the region near a local minimum. Moreover, [55]

proved that the cost function of the policy optimization for the linear quadratic regulator problem is nonconvex and satisfies the P-L condition. More examples of nonconvex functions satisfying the P-L condition can be found in [52], [53].

Although it is difficult to precisely characterize the general class of functions for which the P-L condition is satisfied, in [52], one important special class was given as follows:

Lemma 1: Let $f(x) = g(Ax)$, where $g: \mathbb{R}^p \rightarrow \mathbb{R}$ is a strongly convex function and $A \in \mathbb{R}^{p \times p}$ is a matrix, then f satisfies the P-L condition.

We have the following global convergence results.

Theorem 2: Suppose Assumptions 1–7 hold. For any given $T \geq (c_0(\kappa_1, \kappa_2)/\kappa_2)^{1/\theta}$, let $\{x_0, \dots, x_T\}$ be the sequence generated by Algorithm 1 with

$$\alpha_k = \kappa_1 \beta_k, \beta_k = \kappa_2 (T+1)^\theta, \eta_k = \frac{\kappa_2}{\beta_k}, \forall k \leq T \quad (14)$$

where $\theta \in (0, 1)$, $\kappa_1 > c_1$, $\kappa_2 \in (0, c_2(\kappa_1))$. Then,

$$\mathbf{E}\left[\frac{1}{n} \sum_{i=1}^n \|x_{i,T} - \bar{x}_T\|^2\right] = \mathcal{O}\left(\frac{1}{T^{2\theta}}\right) \quad (15a)$$

$$\mathbf{E}[f(\bar{x}_T) - f^*] = \mathcal{O}\left(\frac{1}{nT^\theta}\right) + \mathcal{O}\left(\frac{1}{T^{2\theta}}\right). \quad (15b)$$

Proof: The explicit expressions of the right-hand sides of (15a) and (15b), and the proof are given in Appendix C. It should be highlighted that the omitted constants in the first term on the right-hand side of (15b) do not depend on any parameters related to the communication network. ■

From Theorem 2, we see that the convergence rate is strictly greater than $\mathcal{O}(1/(nT))$. In the following we show that the linear speedup convergence rate $\mathcal{O}(1/(nT))$ can be achieved if the P-L constant ν is known in advance and each $f_i^* > -\infty$, where $f_i^* = \min_{x \in \mathbb{R}^p} f_i(x)$. The total number of iterations T is not needed.

Theorem 3 (Linear Speedup): Suppose Assumptions 1–7 hold, and the P-L constant ν is known in advance, and each $f_i^* > -\infty$. Let $\{x_k\}$ be the sequence generated by Algorithm 1 with

$$\alpha_k = \kappa_1 \beta_k, \beta_k = \kappa_0 (k+t_1), \eta_k = \frac{\kappa_2}{\beta_k}, \forall k \in \mathbb{N}_0 \quad (16)$$

where $\kappa_0 \in [\hat{c}_0 \nu \kappa_2 / 4, \nu \kappa_2 / 4]$, $\kappa_1 > c_1$, $\kappa_2 \in (0, \hat{c}_2(\kappa_1))$, and $t_1 > \hat{c}_3(\kappa_0, \kappa_1, \kappa_2)$ with $\hat{c}_0 \in (0, 1)$ being a constant, $\hat{c}_2(\kappa_1)$ and $\hat{c}_3(\kappa_0, \kappa_1, \kappa_2)$ defined in Appendix D. Then, for any $T \in \mathbb{N}_+$,

$$\mathbf{E}\left[\frac{1}{n} \sum_{i=1}^n \|x_{i,T} - \bar{x}_T\|^2\right] = \mathcal{O}\left(\frac{1}{T^2}\right) \quad (17a)$$

$$\mathbf{E}[f(\bar{x}_T) - f^*] = \mathcal{O}\left(\frac{1}{nT}\right) + \mathcal{O}\left(\frac{1}{T^2}\right). \quad (17b)$$

Proof: The explicit expressions of the right-hand sides of (17a) and (17b), and the proof are given in Appendix D. It should be highlighted that the omitted constants in the first term on the right-hand side of (17b) do not depend on any parameters related to the communication network. ■

Remark 3: It has been shown in [51] that $\mathcal{O}(1/T)$ convergence rate is optimal for centralized strongly convex optimi-

zation. This rate has been established by various distributed SGD algorithms when each local cost function is strongly convex, e.g., [18], [26], [42]–[46]. In contrast, the linear speedup convergence rate $\mathcal{O}(1/(nT))$ established in Theorem 3 only requires that the global cost function satisfies the P-L condition, but no convexity assumption is required neither on the global cost function nor on the local cost functions. The SGD algorithms in [24], [25], [34], [39]–[41] also achieve the same linear speedup convergence rate. However, the algorithms in [24], [25], [39] are restricted to a star graph, while our algorithm is applicable to an arbitrarily connected graph. Moreover, [24], [25] assumed that the mean of each local stochastic gradient is the gradient of the global cost function, and T has to be known to choose the algorithm parameters. The algorithm in [25] furthermore requires an exponentially increasing batch size, which is not favorable in practice. In [39], it was assumed that the global cost function is strongly convex. In [34], [39], it was assumed that each local stochastic gradient has bounded second moment. In [34], [40], [41], it was assumed that each local cost function is strongly convex. It is one of our future research directions to achieve linear speedup with reduced communication rounds and communication efficiency for an arbitrarily connected graph.

Theorem 3 shows that the convergence rate to a global optimum is sublinear when we allow the parameters to be time-varying. The following theorem establishes that the output of Algorithm 1 with constant algorithm parameters linearly converges to a neighborhood of a global optimum.

Theorem 4: Suppose Assumptions 1–7 hold. Let $\{x_k\}$ be the sequence generated by Algorithm 1 with

$$\alpha_k = \alpha = \kappa_1 \beta, \beta_k = \beta, \eta_k = \eta = \frac{\kappa_2}{\beta}, \forall k \in \mathbb{N}_0 \quad (18)$$

where $\kappa_1 > c_1$, $\kappa_2 \in (0, c_2(\kappa_1))$, and $\beta \geq c_0(\kappa_1, \kappa_2)$ with $c_0(\kappa_1, \kappa_2)$, c_1 , $c_2(\kappa_1) > 0$ defined in Appendix B. Then,

$$\begin{aligned} \mathbf{E}\left[\frac{1}{n} \sum_{i=1}^n \|x_{i,k} - \bar{x}_k\|^2 + f(\bar{x}_k) - f^*\right] \\ \leq (1 - \eta\varepsilon)^k c_4 + c_5 \eta \sigma^2, \forall k \in \mathbb{N}_+ \end{aligned} \quad (19)$$

where $\varepsilon \in (0, 1/\eta)$, $c_4, c_5 > 0$ are constants defined in Appendix E.

Proof: The proof is given in Appendix E. ■

Remark 4: It should be highlighted the P-L constant ν is not used to design the algorithm parameters. Therefore, the constant ν does not need to be known in advance. Similar convergence result as stated in (19) was achieved by the distributed SGD algorithms proposed in [26], [46]–[49] when each local cost function is strongly convex, which obviously is stronger than the P-L condition assumed in Theorem 4. In addition to the strong convexity condition, in [26], it was also assumed that each local cost function is Lipschitz-continuous. Some information related to the Lyapunov function and global parameters, which may be difficult to get, were furthermore needed to design the stepsize. Moreover, in [46]–[49], the strong convexity constant was needed to design the stepsize and in [48], [49], a p -dimensional auxiliary variable, which is used to track the global gradient, was communicated between

SGD-T and DPD-SGD-F) with state-of-the-art algorithms: distributed momentum SGD algorithm (DM-SGD) [23], distributed SGD algorithm (D-SGD-1) [26], [27], distributed SGD algorithm (D-SGD-2) [28], D² [36], distributed stochastic gradient tracking algorithm (D-SGT-1) [37], [49], distributed stochastic gradient tracking algorithm (D-SGT-2) [38], [48], and the baseline centralized SGD algorithm (C-SGD). We list all the parameters¹ we choose in the NN experiment for each algorithm in Table II.

TABLE II
PARAMETERS IN EACH ALGORITHM IN NN EXPERIMENT

Algorithm	η_k	α_k	β_k
DPD-SGD-T	$0.08/k^{10^{-5}}$	$4k^{10^{-5}}$	$3k^{10^{-5}}$
DPD-SGD-F	0.03	5	20
DM-SGD [23]	0.1	×	0.8
D-SGD-1 [26], [27]	0.1	×	×
D-SGD-2 [28]	×	$0.1/(10^{-5}k+1)$	$0.2/(10^{-5}k+1)^{0.3}$
D ² [36]	0.01	×	×
D-SGT-1 [37], [49]	0.01	×	×
D-SGT-2 [38], [48]	0.01	×	×
C-SGD	0.1	×	×

We demonstrate the result in terms of the empirical risk function [60], which is given as

$$R(\mathbf{z}) = -\frac{1}{n} \sum_{i=1}^n \frac{1}{m_n} \sum_{j=1}^{m_n} \sum_{k=0}^9 (t_k \ln y_k(\mathbf{x}, \mathbf{z}) + (1-t_k) \ln(1-y_k(\mathbf{x}, \mathbf{z})))$$

where m_n indicates the size of data set for each agent, t_k denotes the target (ground truth) of digit k corresponding to a single image, \mathbf{x} is a single image input, $\mathbf{z} = (z^{(1)}, z^{(2)})$ with $z^{(1)}$ and $z^{(2)}$ being the weights in the 2 layers respectively, and $y_k \in [0, 1]$ is the output which expresses the probability of digit $k = 0, \dots, 9$. The mapping from input to output is given as

$$y_k(\mathbf{x}, \mathbf{z}) = \sigma \left(\sum_{j=0}^{50} z_{k,j}^{(2)} \sigma \left(\sum_{i=0}^{28 \times 28} z_{j,i}^{(1)} x_i \right) \right)$$

where $\sigma(s) = \frac{1}{1+\exp(-s)}$ is the sigmoid function.

Fig. 2 shows that the proposed distributed primal-dual SGD algorithms with time-varying parameters converges almost as fast as the distributed SGD algorithm in [26], [27] and faster than the distributed SGD algorithms in [28], [36]–[38], [48], [49] and the centralized SGD algorithm. Note that our algorithm converges slower than the distributed momentum SGD algorithm [23]. This is reasonable since that algorithm is an accelerated algorithm with extra requirement on the cost functions, i.e., the deviations between the gradients of local cost functions are bounded, and it requires each agent to communicate three p -dimensional variables with its neighbors at each iteration. The slopes of the curves are however almost the same. The accuracy of each algorithm is given in Table III.

¹ Note: the parameter names are different in each paper.




Fig. 2. Empirical risk.

TABLE III
ACCURACY ON EACH ALGORITHM IN NN EXPERIMENT

Algorithm	Accuracy
DPD-SGD-T	93.04%
DPD-SGD-F	92.76%
DM-SGD [23]	93.44%
D-SGD-1 [26], [27]	92.96%
D-SGD-2 [28]	92.88%
D ² [36]	90.44%
D-SGT-1 [37], [49]	92.88%
D-SGT-2 [38], [48]	92.96%
C-SGD	93%

B. Convolutional Neural Networks

Let us consider the training of a convolutional neural networks (CNN) model. We build a CNN model for each agent with five 3×3 convolutional layers using ReLU as activation function, one average pooling layer with filters of size 2×2 , one sigmoid layer with dimension 360, another sigmoid layer with dimension 60, one softmax layer with dimension 10. In this experiment, we use the whole MNIST data set. We use the same communication graph as in the above NN experiment. Each agent is assigned 6000 data points randomly. We set the batch size as 20, which means at each iteration, 20 data points are chosen by the agent to update the gradient, which is also following a uniform distribution. For each algorithm, we do 10 epochs to train the CNN model.

We compare our algorithms DPD-SGD-T and DPD-SGD-F with the fastest ones for the neural networks case, i.e., DM-SGD [23], D-SGD-1 [26], [27], and C-SGD. We list all the parameters we choose in the CNN experiment for each algorithm in Table IV.

We demonstrate the training loss and the test accuracy of each algorithm in Figs. 3 and 4 respectively. Here we use Categorical Cross-Entropy loss, which is a softmax activation plus a Cross-Entropy loss. We can see that our algorithms perform almost the same as the DM-SGD and better than the D-SGD-1 and the centralized C-SGD. The accuracy of each algorithm is given in Table V.

TABLE IV
PARAMETERS IN EACH ALGORITHM IN CNN EXPERIMENT

Algorithm	η_k	α_k	β_k
DPD-SGD-T	$0.5/k^{10^{-5}}$	$0.5k^{10^{-5}}$	$0.1k^{10^{-5}}$
DPD-SGD-F	0.5	0.5	0.1
DM-SGD [23]	0.1	×	0.8
D-SGD-1 [26], [27]	0.1	×	×
C-SGD	0.1	×	×




Fig. 3. CNN training loss.




Fig. 4. CNN accuracy.

TABLE V
ACCURACY ON EACH ALGORITHM IN CNN EXPERIMENT

Algorithm	Accuracy
DPD-SGD-T	94.75%
DPD-SGD-F	93.17%
DM-SGD [23]	94.29%
D-SGD-1 [26], [27]	92.96%
C-SGD	89.91%

V. CONCLUSIONS

In this paper, we studied distributed nonconvex optimization. We proposed a distributed primal-dual SGD algorithm and derived its convergence rate. More specifically, the linear

speedup convergence rate $O(1/\sqrt{nT})$ was established for smooth nonconvex cost functions under arbitrarily connected communication networks. The convergence rate was improved to the linear speedup convergence rate $O(1/(nT))$ when the global cost function additionally satisfies the P-L condition. It was also shown that the output of the proposed algorithm with constant parameters linearly converges to a neighborhood of a global optimum. Interesting directions for future work include achieving linear speedup under the P-L condition while considering communication reduction.

APPENDIX

A. Notations and Useful Lemmas

\mathbf{I}_n is the n -dimensional identity matrix. The notation $A \otimes B$ denotes the Kronecker product of matrices A and B . $\text{null}(A)$ is the null space of matrix A . Given two symmetric matrices M, N , $M \geq N$ means that $M - N$ is positive semi-definite. $\rho(\cdot)$ stands for the spectral radius of a matrix and $\rho_2(\cdot)$ indicates the minimum positive eigenvalue for a matrix having positive eigenvalues. For any square matrix A , $\|x\|_A^2$ denotes $x^T A x$. $\lceil \cdot \rceil$ and $\lfloor \cdot \rfloor$ denote the ceiling and floor functions, respectively. For any $x \in \mathbb{R}$, $[x]_+$ is the positive part of x . $\mathbf{1}_{(\cdot)}$ is the indicator function. For any $n \in \mathbb{N}_0$, $n!$ is the factorial of n .

Denote $K_n = \mathbf{I}_n - \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T$, $\mathbf{K} = K_n \otimes \mathbf{I}_p$, $\mathbf{H} = \frac{1}{n} (\mathbf{1}_n \mathbf{1}_n^T \otimes \mathbf{I}_p)$, $\bar{x}_k = \frac{1}{n} (\mathbf{1}_n^T \otimes \mathbf{I}_p) \mathbf{x}_k$, $\bar{x}_k = \mathbf{1}_n \otimes \bar{x}_k$, $\mathbf{g}_k = \nabla \tilde{f}(\mathbf{x}_k)$, $\bar{\mathbf{g}}_k = \mathbf{H} \mathbf{g}_k$, $\mathbf{g}_k^0 = \nabla \tilde{f}(\bar{x}_k)$, $\bar{\mathbf{g}}_k^0 = \mathbf{H} \mathbf{g}_k^0 = \mathbf{1}_n \otimes \nabla f(\bar{x}_k)$, and $\bar{\mathbf{g}}_k^\mu = \mathbf{H} \mathbf{g}_k^\mu$.

The following results are used in the proofs.

Lemma 2 (Lemma 1.2.3 in [61] and Lemma 3 in [62]): If the function $f(x) : \mathbb{R}^p \mapsto \mathbb{R}$ is smooth with constant $L_f > 0$, then,

$$|f(y) - f(x) - (y-x)^T \nabla f(x)| \leq \frac{L_f}{2} \|y-x\|^2 \quad (24a)$$

$$\|\nabla f(x)\|^2 \leq 2L_f(f(x) - f^*), \quad \forall x, y \in \mathbb{R}^p \quad (24b)$$

where $f^* = \min_{x \in \mathbb{R}^p} f(x)$.

Lemma 3 (Lemmas 1 and 2 in [63]): Let L be the Laplacian matrix of the graph \mathcal{G} . If Assumption 1 holds, then L is positive semi-definite, $\text{null}(L) = \text{null}(K_n) = \{\mathbf{1}_n\}$, $L \leq \rho(L) \mathbf{I}_n$, $\rho(K_n) = 1$,

$$K_n L = L K_n = L \quad (25)$$

$$0 \leq \rho_2(L) K_n \leq L \leq \rho(L) K_n. \quad (26)$$

Moreover, there exists an orthogonal matrix $[r \ R] \in \mathbb{R}^{n \times n}$ with $r = \frac{1}{\sqrt{n}} \mathbf{1}_n$ and $R \in \mathbb{R}^{n \times (n-1)}$ such that

$$R \Lambda_1^{-1} R^T L = L R \Lambda_1^{-1} R^T = K_n \quad (27)$$

$$\frac{1}{\rho(L)} K_n \leq R \Lambda_1^{-1} R^T \leq \frac{1}{\rho_2(L)} K_n \quad (28)$$

where $\Lambda_1 = \text{diag}([\lambda_2, \dots, \lambda_n])$ with $0 < \lambda_2 \leq \dots \leq \lambda_n$ being the eigenvalues of the Laplacian matrix L .

Lemma 4: Let $a \in (0, 1)$ be a constant, then

$$(1-a)^T \leq \frac{k!}{(aT)^k}, \quad \forall k, T \in \mathbb{N}_0. \quad (29)$$

Proof: For any constant $a \in (0, 1)$, we have $\ln(1-a) \leq -a$. Thus,

$$(1-a)^T \leq e^{-aT}, \forall T \in \mathbb{N}_0. \quad (30)$$

For any constant $x > 0$, we have $e^x > \frac{x^k}{k!}$, $\forall k \in \mathbb{N}_0$. This result together with (30) yields (29). ■

Lemma 5: Let $\{z_k\}$, $\{r_{1,k}\}$, and $\{r_{2,k}\}$ be sequences. Suppose there exists $t_1 \in \mathbb{N}_+$ such that

$$z_k \geq 0 \quad (31a)$$

$$z_{k+1} \leq (1-r_{1,k})z_k + r_{2,k} \quad (31b)$$

$$1 > r_{1,k} \geq \frac{a_1}{(k+t_1)^\delta} \quad (31c)$$

$$r_{2,k} \leq \frac{a_2}{(k+t_1)^2}, \forall k \in \mathbb{N}_0 \quad (31d)$$

where $\delta \geq 0$, $a_1 > 0$, and $a_2 > 0$ are constants.

i) If $\delta = 1$, then,

$$z_k \leq \phi_1(k, t_1, a_1, a_2, z_0), \forall k \in \mathbb{N}_+ \quad (32)$$

where

$$\phi_1(k, t_1, a_1, a_2, z_0) = \frac{t_1^{a_1} z_0}{(k+t_1)^{a_1}} + \frac{a_2}{(k+t_1-1)^2} + 4a_2 s_1(k+t_1) \quad (33)$$

and

$$s_1(k) = \begin{cases} \frac{1}{(a_1-1)k}, & \text{if } a_1 > 1 \\ \frac{\ln(k-1)}{k}, & \text{if } a_1 = 1 \\ \frac{-t_1^{a_1-1}}{(a_1-1)k^{a_1}}, & \text{if } a_1 < 1. \end{cases}$$

ii) If $\delta = 0$, then,

$$z_k \leq \phi_2(k, t_1, a_1, a_2, z_0), \forall k \in \mathbb{N}_+ \quad (34)$$

where

$$\begin{aligned} \phi_2(k, t_1, a_1, a_2, z_0) &= (1-a_1)^k z_0 + a_2 (1-a_1)^{k+t_1-1} \left([t_2-t_1]_+ s_2(t_1) \right. \\ &\quad \left. + ([t_3-t_1]_+ - [t_2-t_1]_+) s_2(t_3) \right) \\ &\quad + \frac{\mathbf{1}_{(k+t_1-1 \geq t_3)} 2a_2}{-\ln(1-a_1)(k+t_1)^2(1-a_1)} \end{aligned} \quad (35)$$

$$s_2(k) = \frac{1}{k^2(1-a_1)^k}, t_2 = \lceil \frac{-2}{\ln(1-a_1)} \rceil, \text{ and } t_3 = \lceil \frac{-4}{\ln(1-a_1)} \rceil.$$

Proof: i) From (31a)–(31c), for any $k \in \mathbb{N}_+$, it holds that

$$z_k \leq \prod_{\tau=0}^{k-1} (1-r_{1,\tau}) z_0 + r_{2,k-1} + \sum_{l=0}^{k-2} \prod_{\tau=l+1}^{k-1} (1-r_{1,\tau}) r_{2,l}. \quad (36)$$

For any $t \in [0, 1]$, it holds that $1-t \leq e^{-t}$ since $s_3(t) = 1-t-e^{-t}$ is a non-increasing function in the interval $[0, 1]$. Thus, for any $k > l \geq 0$, it holds that

$$\prod_{\tau=l}^{k-1} (1-r_{1,\tau}) \leq e^{-\sum_{\tau=l}^{k-1} r_{1,\tau}}. \quad (37)$$

We also have

$$\begin{aligned} \sum_{\tau=l}^{k-1} r_{1,\tau} &\geq \sum_{\tau=l}^{k-1} \frac{a_1}{\tau+t_1} = \sum_{\tau=l+t_1}^{k-1+t_1} \frac{a_1}{\tau} \\ &\geq \int_{l+t_1}^{k+t_1} \frac{a_1}{t} dt = a_1 (\ln(k+t_1) - \ln(l+t_1)) \end{aligned} \quad (38)$$

where the first inequality holds due to (31c) and the second inequality holds since $s_4(t) = a_1/t$ is a decreasing function in the interval $[1, +\infty)$.

Hence, (37) and (38) yield

$$\prod_{\tau=l}^{k-1} (1-r_{1,\tau}) \leq e^{-\sum_{\tau=l}^{k-1} r_{1,\tau}} \leq \frac{(l+t_1)^{a_1}}{(k+t_1)^{a_1}}. \quad (39)$$

We have

$$\begin{aligned} \sum_{l=0}^{k-2} \prod_{\tau=l+1}^{k-1} (1-r_{1,\tau}) r_{2,l} &\leq \sum_{l=0}^{k-2} \frac{(l+t_1+1)^{a_1}}{(k+t_1)^{a_1}} \frac{a_2}{(l+t_1)^2} \\ &\leq \sum_{l=0}^{k-2} \frac{(l+t_1+1)^{a_1}}{(k+t_1)^{a_1}} \frac{a_2}{(\frac{t_1}{l+1}l+t_1)^2} \\ &= \frac{(\frac{t_1+1}{t_1})^2 a_2}{(k+t_1)^{a_1}} \sum_{l=0}^{k-2} \frac{(l+t_1+1)^{a_1}}{(l+t_1+1)^2} \\ &= \frac{4a_2}{(k+t_1)^{a_1}} \sum_{l=t_1+1}^{k+t_1-1} l^{a_1-2} \end{aligned} \quad (40)$$

where the first inequality holds due to (39) and (31d).

From (36), (39), and (40), we have (32).

ii) Denote $a = 1-a_1$. From (31c) and $\delta = 0$, we know that $a_1 \in (0, 1)$. Thus, $a \in (0, 1)$.

From (31a)–(31d) and $\delta_1 = 0$, for any $k \in \mathbb{N}_+$, it holds that

$$\begin{aligned} z_k &\leq (1-a_1)^k z_0 + \sum_{\tau=0}^{k-1} (1-a_1)^{k-1-\tau} r_{2,\tau} \\ &\leq a^k z_0 + a_2 a^{k+t_1-1} \sum_{\tau=0}^{k-1} \frac{1}{(\tau+t_1)^2 a^{\tau+t_1}}. \end{aligned} \quad (41)$$

We have

$$\begin{aligned} \sum_{\tau=0}^{k-1} \frac{1}{(\tau+t_1)^2 a^{\tau+t_1}} &= \sum_{\tau=t_1}^{k+t_1-1} \frac{1}{\tau^2 a^\tau} \\ &= \sum_{\tau=t_1}^{t_2-1} s_2(\tau) + \sum_{\tau=t_2}^{t_3-1} s_2(\tau) + \sum_{\tau=t_3}^{k+t_1-1} s_2(\tau). \end{aligned} \quad (42)$$

We know that $s_2(t) = 1/(t^2 a^t)$ is decreasing and increasing in the intervals $[1, t_2-1]$ and $[t_2, +\infty)$, respectively, since

$$\frac{ds_2(t)}{dt} = -s_2(t) \left(\frac{2}{t} + \ln(a) \right) \leq 0, \forall t \in \left(0, \frac{-2}{\ln(a)} \right],$$

$$\frac{ds_2(t)}{dt} = -s_2(t) \left(\frac{2}{t} + \ln(a) \right) \geq 0, \forall t \in \left[\frac{-2}{\ln(a)}, +\infty \right).$$

Thus, we have

$$\sum_{\tau=k_1}^{t_2-1} s_2(\tau) \leq (t_2-k_1) s_2(k_1), \forall k_1 \in [1, t_2-1] \quad (43a)$$

$$\sum_{\tau=k_2}^{t_3-1} s_2(\tau) \leq (t_3 - k_2)s_2(t_3), \quad \forall k_2 \in [t_2, t_3 - 1] \quad (43b)$$

$$\sum_{\tau=t_3}^{k_3} s_2(\tau) \leq \int_{t_3}^{k_3+1} s_2(t)dt, \quad \forall k_3 \geq t_3. \quad (43c)$$

Denote $b = 1/a$. We have

$$\begin{aligned} \int_{t_3}^{k_3+1} s_2(t)dt &= \int_{t_3}^{k_3+1} \frac{b^t}{t^2} dt = \int_{t_3}^{k_3+1} \frac{1}{\ln(b)t^2} db^t \\ &= \frac{b^{k_3+1}}{\ln(b)(k_3+1)^2} - \frac{b^{t_3}}{\ln(b)t_3^2} + \int_{t_3}^{k_3+1} \frac{2b^t}{\ln(b)t^3} dt \\ &\leq \frac{b^{k_3+1}}{\ln(b)(k_3+1)^2} + \int_{t_3}^{k_3+1} \frac{2}{\ln(b)t} s_2(t)dt \\ &\leq \frac{b^{k_3+1}}{\ln(b)(k_3+1)^2} + \frac{2}{\ln(b)t_3} \int_{t_3}^{k_3+1} s_2(t)dt \\ &\leq \frac{b^{k_3+1}}{\ln(b)(k_3+1)^2} + \frac{1}{2} \int_{t_3}^{k_3+1} s_2(t)dt \end{aligned} \quad (44)$$

where the last inequality holds due to $t_3 = \lceil \frac{-4}{\ln(1-a_1)} \rceil \geq \frac{-4}{\ln(1-a_1)} = \frac{4}{\ln(b)}$.

From (43c) and (44), we have

$$\sum_{\tau=t_3}^{k_3} s_2(\tau) \leq \frac{-2}{\ln(a)(k_3+1)^2 a^{k_3+1}}, \quad \forall k_3 \geq t_3. \quad (45)$$

From (41), (42), (43a), (43b), and (45), we get (34). ■

Lemma 6: Suppose Assumptions 1 and 3–6 hold. Then the following holds for Algorithm 1:

$$\begin{aligned} \mathbf{E}_{\tilde{\delta}_k} [W_{1,k+1}] &\leq W_{1,k} - \|\mathbf{x}_k\|_{\eta_k \alpha_k L - \frac{1}{2} \eta_k \mathbf{K} - \frac{3}{2} \eta_k^2 \alpha_k^2 L^2 - \frac{1}{2} \eta_k (1+5\eta_k) L_f^2 \mathbf{K}} \\ &\quad - \eta_k \beta_k \mathbf{x}_k^T \mathbf{K} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) + \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\frac{3}{2} \eta_k^2 \beta_k^2 \mathbf{K}} \\ &\quad + 2n\sigma^2 \eta_k^2 \end{aligned} \quad (46)$$

where $W_{1,k} = \frac{1}{2} \|\mathbf{x}_k\|_{\mathbf{K}}^2$.

Proof: Noting that $\nabla \tilde{f}$ is Lipschitz-continuous with constant $L_f > 0$ since Assumption 3 is satisfied, we have that

$$\|\mathbf{g}_k^0 - \mathbf{g}_k\|^2 \leq L_f^2 \|\bar{\mathbf{x}}_k - \mathbf{x}_k\|^2 = L_f^2 \|\mathbf{x}_k\|_{\mathbf{K}}^2. \quad (47)$$

From Assumptions 4–6, we know that

$$\mathbf{E}_{\tilde{\delta}_k} [\mathbf{g}_k^u] = \mathbf{g}_k \quad (48a)$$

$$\mathbf{E}_{\tilde{\delta}_k} [\|\mathbf{g}_k^u - \mathbf{g}_k\|^2] \leq n\sigma^2. \quad (48b)$$

From (47), (48b), and the Cauchy-Schwarz inequality, we have

$$\begin{aligned} \mathbf{E}_{\tilde{\delta}_k} [\|\mathbf{g}_k^0 - \mathbf{g}_k^u\|^2] &= \mathbf{E}_{\tilde{\delta}_k} [\|\mathbf{g}_k^0 - \mathbf{g}_k + \mathbf{g}_k - \mathbf{g}_k^u\|^2] \\ &\leq 2\|\mathbf{g}_k^0 - \mathbf{g}_k\|^2 + 2\mathbf{E}_{\tilde{\delta}_k} [\|\mathbf{g}_k - \mathbf{g}_k^u\|^2] \\ &\leq 2L_f^2 \|\mathbf{x}_k\|_{\mathbf{K}}^2 + 2n\sigma^2. \end{aligned} \quad (49)$$

We have

$$\begin{aligned} \mathbf{E}_{\tilde{\delta}_k} [W_{1,k+1}] &= \mathbf{E}_{\tilde{\delta}_k} \left[\frac{1}{2} \|\mathbf{x}_{k+1}\|_{\mathbf{K}}^2 \right] \\ &= \mathbf{E}_{\tilde{\delta}_k} \left[\frac{1}{2} \|\mathbf{x}_k - \eta_k (\alpha_k \mathbf{L} \mathbf{x}_k + \beta_k \mathbf{v}_k + \mathbf{g}_k^u)\|_{\mathbf{K}}^2 \right] \\ &= \mathbf{E}_{\tilde{\delta}_k} \left[\frac{1}{2} \|\mathbf{x}_k\|_{\mathbf{K}}^2 - \eta_k \alpha_k \|\mathbf{x}_k\|_{\mathbf{L}}^2 + \frac{1}{2} \eta_k^2 \alpha_k^2 \|\mathbf{x}_k\|_{\mathbf{L}^2}^2 \right. \\ &\quad \left. - \eta_k \beta_k \mathbf{x}_k^T (\mathbf{I}_{np} - \eta_k \alpha_k \mathbf{L}) \mathbf{K} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^u \right) \right. \\ &\quad \left. + \frac{1}{2} \eta_k^2 \beta_k^2 \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^u \right\|_{\mathbf{K}}^2 \right] \\ &= \frac{1}{2} \|\mathbf{x}_k\|_{\mathbf{K}}^2 - \|\mathbf{x}_k\|_{\eta_k \alpha_k L - \frac{1}{2} \eta_k^2 \alpha_k^2 L^2}^2 - \eta_k \beta_k \mathbf{x}_k^T (\mathbf{I}_{np} \\ &\quad - \eta_k \alpha_k \mathbf{L}) \mathbf{K} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 + \frac{1}{\beta_k} \mathbf{g}_k - \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \\ &\quad + \frac{1}{2} \eta_k^2 \beta_k^2 \mathbf{E}_{\tilde{\delta}_k} \left[\left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 + \frac{1}{\beta_k} \mathbf{g}_k^u - \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\mathbf{K}}^2 \right] \\ &\leq W_{1,k} - \|\mathbf{x}_k\|_{\eta_k \alpha_k L - \frac{1}{2} \eta_k^2 \alpha_k^2 L^2}^2 \\ &\quad - \eta_k \beta_k \mathbf{x}_k^T \mathbf{K} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \\ &\quad + \frac{\eta_k}{2} \|\mathbf{x}_k\|_{\mathbf{K}}^2 + \frac{\eta_k}{2} \|\mathbf{g}_k - \mathbf{g}_k^0\|^2 \\ &\quad + \frac{1}{2} \eta_k^2 \alpha_k^2 \|\mathbf{x}_k\|_{\mathbf{L}^2}^2 + \frac{1}{2} \eta_k^2 \beta_k^2 \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\mathbf{K}}^2 \\ &\quad + \frac{1}{2} \eta_k^2 \alpha_k^2 \|\mathbf{x}_k\|_{\mathbf{L}^2}^2 + \frac{1}{2} \eta_k^2 \|\mathbf{g}_k - \mathbf{g}_k^0\|^2 \\ &\quad + \eta_k^2 \beta_k^2 \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\mathbf{K}}^2 + \eta_k^2 \mathbf{E}_{\tilde{\delta}_k} [\|\mathbf{g}_k^u - \mathbf{g}_k^0\|^2] \\ &= W_{1,k} - \|\mathbf{x}_k\|_{\eta_k \alpha_k L - \frac{1}{2} \eta_k \mathbf{K} - \frac{3}{2} \eta_k^2 \alpha_k^2 L^2}^2 \\ &\quad + \frac{\eta_k}{2} (1 + \eta_k) \|\mathbf{g}_k - \mathbf{g}_k^0\|^2 + \eta_k^2 \mathbf{E}_{\tilde{\delta}_k} [\|\mathbf{g}_k^u - \mathbf{g}_k^0\|^2] \\ &\quad - \eta_k \beta_k \mathbf{x}_k^T \mathbf{K} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) + \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\frac{3}{2} \eta_k^2 \beta_k^2 \mathbf{K}}^2 \end{aligned} \quad (50)$$

where the second equality holds due to (5a); the third equality holds due to (25) in Lemma 3; the fourth equality holds since \mathbf{x}_k and \mathbf{v}_k are independent of $\tilde{\delta}_k$ and (48a), respectively; and the inequality holds due to the Cauchy-Schwarz inequality and $\rho(\mathbf{K}) = 1$.

Then, from (47), (49), and (50), we have (46). ■

Lemma 7: Suppose Assumptions 1 and 3 hold, and $\{\beta_k\}$ is non-decreasing. Then the following holds for Algorithm 1:

$$\begin{aligned} W_{2,k+1} &\leq W_{2,k} + (1 + \omega_k) \eta_k \beta_k \mathbf{x}_k^T (\mathbf{K} + \kappa_1 \mathbf{L}) \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \\ &\quad + \frac{1}{2} (\eta_k + \omega_k + \eta_k \omega_k) \left(\frac{1}{\rho_2(L)} + \kappa_1 \right) \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\mathbf{K}}^2 \\ &\quad + \|\mathbf{x}_k\|_{(1+\omega_k) \eta_k^2 \beta_k^2 (\mathbf{L} + \kappa_1 \mathbf{L}^2)}^2 \\ &\quad + \frac{\eta_k}{\beta_k^2} \left(\eta_k + \frac{1}{2} \right) (1 + \omega_k) \left(\frac{1}{\rho_2(L)} + \kappa_1 \right) L_f^2 \|\bar{\mathbf{g}}_k^u\|^2 \\ &\quad + \frac{1}{2} \left(\frac{1}{\rho_2(L)} + \kappa_1 \right) (\omega_k + \omega_k^2) \|\mathbf{g}_{k+1}^0\|^2 \end{aligned} \quad (51)$$

where $W_{2,k} = \frac{1}{2} \|\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2$, $\mathbf{Q} = R\Lambda_1^{-1}R^\top \otimes \mathbf{I}_p$ with matrices R and Λ_1^{-1} given in Lemma 3, $\omega_k = \frac{1}{\beta_k} - \frac{1}{\beta_{k+1}}$, and $\kappa_1 > 0$ is a constant.

Proof: Denote $\bar{\mathbf{v}}_k = \frac{1}{n}(\mathbf{1}_n^\top \otimes \mathbf{I}_p)\mathbf{v}_k$. Then, from (5b), we know that

$$\bar{\mathbf{v}}_{k+1} = \bar{\mathbf{v}}_k. \quad (52)$$

Then, from (52) and $\sum_{i=1}^n v_{i,0} = \mathbf{0}_p$, it can be obtained that

$$\bar{\mathbf{v}}_k = \mathbf{0}_p. \quad (53)$$

Then, from (53) and (5a), we have

$$\bar{\mathbf{x}}_{k+1} = \bar{\mathbf{x}}_k - \eta_k \bar{\mathbf{g}}_k^u. \quad (54)$$

Since $\nabla \tilde{f}$ is Lipschitz-continuous and (54), we have

$$\|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0\|^2 \leq L_f^2 \|\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}_k\|^2 = \eta_k^2 L_f^2 \|\bar{\mathbf{g}}_k^u\|^2. \quad (55)$$

We know that $\omega_k \geq 0$ since $\{\beta_k\}$ is non-decreasing. We have

$$\begin{aligned} W_{2,k+1} &= \frac{1}{2} \left\| \mathbf{v}_{k+1} + \frac{1}{\beta_{k+1}} \mathbf{g}_{k+1}^0 \right\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \\ &= \frac{1}{2} \left\| \mathbf{v}_{k+1} + \frac{1}{\beta_k} \mathbf{g}_{k+1}^0 + \left(\frac{1}{\beta_{k+1}} - \frac{1}{\beta_k} \right) \mathbf{g}_{k+1}^0 \right\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \\ &\leq \frac{1}{2} (1 + \omega_k) \left\| \mathbf{v}_{k+1} + \frac{1}{\beta_k} \mathbf{g}_{k+1}^0 \right\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \\ &\quad + \frac{1}{2} (\omega_k + \omega_k^2) \|\mathbf{g}_{k+1}^0\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \end{aligned} \quad (56)$$

where the inequality holds due to the Cauchy-Schwarz inequality.

For the first term on the right-hand side of (56), we have

$$\begin{aligned} &\frac{1}{2} \left\| \mathbf{v}_{k+1} + \frac{1}{\beta_k} \mathbf{g}_{k+1}^0 \right\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \\ &= \frac{1}{2} \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 + \eta_k \beta_k \mathbf{L} \mathbf{x}_k + \frac{1}{\beta_k} (\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0) \right\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \\ &= \frac{1}{2} \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \\ &\quad + \eta_k \beta_k \mathbf{x}_k^\top (\mathbf{K} + \kappa_1 \mathbf{L}) \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \\ &\quad + \|\mathbf{x}_k\|_{\frac{1}{2} \eta_k^2 \beta_k^2 (\mathbf{L} + \kappa_1 \mathbf{L}^2)}^2 + \frac{1}{2\beta_k^2} \|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \\ &\quad + \frac{1}{\beta_k} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 + \eta_k \beta_k \mathbf{L} \mathbf{x}_k \right)^\top (\mathbf{Q} + \kappa_1 \mathbf{K}) (\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0) \\ &\leq W_{2,k} + \eta_k \beta_k \mathbf{x}_k^\top (\mathbf{K} + \kappa_1 \mathbf{L}) \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \\ &\quad + \|\mathbf{x}_k\|_{\frac{1}{2} \eta_k^2 \beta_k^2 (\mathbf{L} + \kappa_1 \mathbf{L}^2)}^2 + \frac{1}{2\beta_k^2} \|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \\ &\quad + \frac{\eta_k}{2} \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 + \frac{1}{2\eta_k \beta_k^2} \|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \\ &\quad + \frac{1}{2} \eta_k^2 \beta_k^2 \|\mathbf{L} \mathbf{x}_k\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 + \frac{1}{2\beta_k^2} \|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \\ &= W_{2,k} + \eta_k \beta_k \mathbf{x}_k^\top (\mathbf{K} + \kappa_1 \mathbf{L}) \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \end{aligned}$$

$$\begin{aligned} &+ \|\mathbf{x}_k\|_{\frac{1}{2} \eta_k^2 \beta_k^2 (\mathbf{L} + \kappa_1 \mathbf{L}^2)}^2 + \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\frac{1}{2} \eta_k (\mathbf{Q} + \kappa_1 \mathbf{K})}^2 \\ &+ \frac{1}{\beta_k^2} \left(1 + \frac{1}{2\eta_k} \right) \|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \\ &\leq W_{2,k} + \eta_k \beta_k \mathbf{x}_k^\top (\mathbf{K} + \kappa_1 \mathbf{L}) \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \\ &\quad + \|\mathbf{x}_k\|_{\frac{1}{2} \eta_k^2 \beta_k^2 (\mathbf{L} + \kappa_1 \mathbf{L}^2)}^2 + \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\frac{1}{2} \eta_k (\mathbf{Q} + \kappa_1 \mathbf{K})}^2 \\ &\quad + \frac{1}{\beta_k^2} \left(1 + \frac{1}{2\eta_k} \right) \left(\frac{1}{\rho_2(L)} + \kappa_1 \right) \|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0\|^2 \\ &\leq W_{2,k} + \eta_k \beta_k \mathbf{x}_k^\top (\mathbf{K} + \kappa_1 \mathbf{L}) \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \\ &\quad + \|\mathbf{x}_k\|_{\frac{1}{2} \eta_k^2 \beta_k^2 (\mathbf{L} + \kappa_1 \mathbf{L}^2)}^2 + \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\frac{1}{2} \eta_k (\mathbf{Q} + \kappa_1 \mathbf{K})}^2 \\ &\quad + \frac{\eta_k}{\beta_k^2} \left(\eta_k + \frac{1}{2} \right) \left(\frac{1}{\rho_2(L)} + \kappa_1 \right) L_f^2 \|\bar{\mathbf{g}}_k^u\|^2 \end{aligned} \quad (57)$$

where the first equality holds due to (5b); the second equality holds due to (25) and (27) in Lemma 3; the first inequality holds due to the Cauchy-Schwarz inequality; the last equality holds due to (25) and (27) in Lemma 3; the second inequality holds due to $\rho(\mathbf{Q} + \kappa_1 \mathbf{K}) \leq \rho(\mathbf{Q}) + \kappa_1 \rho(\mathbf{K})$, (28), $\rho(\mathbf{K}) = 1$; and the last inequality holds due to (55).

For the second term on the right-hand side of (56), we have

$$\|\mathbf{g}_{k+1}^0\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \leq \left(\frac{1}{\rho_2(L)} + \kappa_1 \right) \|\mathbf{g}_{k+1}^0\|^2. \quad (58)$$

Also note that

$$\left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\mathbf{Q} + \kappa_1 \mathbf{K}}^2 \leq \left(\frac{1}{\rho_2(L)} + \kappa_1 \right) \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\mathbf{K}}^2. \quad (59)$$

Then, from (56)–(59), we have (51). \blacksquare

Lemma 8: Suppose Assumptions 1 and 3–6 hold, and $\{\beta_k\}$ is non-decreasing. Then the following holds for Algorithm 1:

$$\begin{aligned} \mathbf{E}_{\bar{\delta}_k} [W_{3,k+1}] &\leq W_{3,k} - (1 + \omega_k) \eta_k \alpha_k \mathbf{x}_k^\top \mathbf{L} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \\ &\quad + \|\mathbf{x}_k\|_{\eta_k (\beta_k \mathbf{L} + \frac{1}{2} \mathbf{K}) + \eta_k^2 (\frac{1}{2} \alpha_k^2 - \alpha_k \beta_k + \beta_k^2) \mathbf{L}^2}^2 \\ &\quad + \|\mathbf{x}_k\|_{\frac{1}{2} \omega_k \eta_k \alpha_k \mathbf{L}^2 + \frac{1}{2} \eta_k (1 + 3\eta_k) \mathbf{L}_f^2 \mathbf{K}}^2 \\ &\quad + \frac{\eta_k}{2\beta_k^2} (1 + 3\eta_k) L_f^2 \mathbf{E}_{\bar{\delta}_k} [\|\bar{\mathbf{g}}_k^u\|^2] + n\sigma^2 \eta_k^2 \\ &\quad - \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\eta_k (\beta_k - \frac{1}{2} - \eta_k \beta_k^2 - \frac{1}{2} \omega_k \alpha_k) \mathbf{K}}^2 \\ &\quad + \frac{1}{2} \omega_k \mathbf{E}_{\bar{\delta}_k} [2W_{1,k+1} + \|\mathbf{g}_{k+1}^0\|^2] \end{aligned} \quad (60)$$

where $W_{3,k} = \mathbf{x}_k^\top \mathbf{K} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right)$.

Proof: We have

$$\begin{aligned} W_{3,k+1} &= \mathbf{x}_{k+1}^\top \mathbf{K} \left(\mathbf{v}_{k+1} + \frac{1}{\beta_{k+1}} \mathbf{g}_{k+1}^0 \right) \\ &= \mathbf{x}_{k+1}^\top \mathbf{K} \left(\mathbf{v}_{k+1} + \frac{1}{\beta_k} \mathbf{g}_{k+1}^0 + \left(\frac{1}{\beta_{k+1}} - \frac{1}{\beta_k} \right) \mathbf{g}_{k+1}^0 \right) \end{aligned}$$

$$\begin{aligned}
&= \mathbf{x}_{k+1}^T \mathbf{K} \left(\mathbf{v}_{k+1} + \frac{1}{\beta_k} \mathbf{g}_{k+1}^0 \right) - \omega_k \mathbf{x}_{k+1}^T \mathbf{K} \mathbf{g}_{k+1}^0 \\
&\leq \mathbf{x}_{k+1}^T \mathbf{K} \left(\mathbf{v}_{k+1} + \frac{1}{\beta_k} \mathbf{g}_{k+1}^0 \right) \\
&\quad + \frac{1}{2} \omega_k (\|\mathbf{x}_{k+1}\|_{\mathbf{K}}^2 + \|\mathbf{g}_{k+1}^0\|^2). \tag{61}
\end{aligned}$$

For the first term on the right-hand side of (61), we have

$$\begin{aligned}
&\mathbf{E}_{\tilde{\delta}_k} \left[\mathbf{x}_{k+1}^T \mathbf{K} \left(\mathbf{v}_{k+1} + \frac{1}{\beta_k} \mathbf{g}_{k+1}^0 \right) \right] \\
&= \mathbf{E}_{\tilde{\delta}_k} \left[(\mathbf{x}_k - \eta_k (\alpha_k \mathbf{L} \mathbf{x}_k + \beta_k \mathbf{v}_k + \mathbf{g}_k^0 + \mathbf{g}_k^\mu - \mathbf{g}_k^0))^T \right. \\
&\quad \times \mathbf{K} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 + \eta_k \beta_k \mathbf{L} \mathbf{x}_k + \frac{1}{\beta_k} (\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0) \right) \left. \right] \\
&= \mathbf{x}_k^T (\mathbf{K} - \eta_k (\alpha_k + \eta_k \beta_k^2) \mathbf{L}) \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \\
&\quad + \|\mathbf{x}_k\|_{\eta_k \beta_k (\mathbf{L} - \eta_k \alpha_k \mathbf{L}^2)}^2 \\
&\quad + \frac{1}{\beta_k} \mathbf{x}_k^T (\mathbf{K} - \eta_k \alpha_k \mathbf{L}) \mathbf{E}_{\tilde{\delta}_k} [\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0] \\
&\quad - \eta_k \beta_k \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\mathbf{K}}^2 \\
&\quad - \eta_k \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right)^T \mathbf{K} \mathbf{E}_{\tilde{\delta}_k} [\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0] \\
&\quad - \eta_k (\mathbf{g}_k - \mathbf{g}_k^0)^T \mathbf{K} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 + \eta_k \beta_k \mathbf{L} \mathbf{x}_k \right) \\
&\quad - \frac{1}{\beta_k} \mathbf{E}_{\tilde{\delta}_k} [\eta_k (\mathbf{g}_k^\mu - \mathbf{g}_k^0)^T \mathbf{K} (\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0)] \\
&\leq \mathbf{x}_k^T (\mathbf{K} - \eta_k \alpha_k \mathbf{L}) \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) + \frac{1}{2} \eta_k^2 \beta_k^2 \|\mathbf{L} \mathbf{x}_k\|^2 \\
&\quad + \frac{1}{2} \eta_k^2 \beta_k^2 \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\mathbf{K}}^2 + \|\mathbf{x}_k\|_{\eta_k \beta_k (\mathbf{L} - \eta_k \alpha_k \mathbf{L}^2)}^2 \\
&\quad + \frac{1}{2} \eta_k \|\mathbf{x}_k\|_{\mathbf{K}}^2 + \frac{1}{2 \eta_k \beta_k^2} \mathbf{E}_{\tilde{\delta}_k} [\|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0\|^2] \\
&\quad + \frac{1}{2} \eta_k^2 \alpha_k^2 \|\mathbf{L} \mathbf{x}_k\|^2 + \frac{1}{2 \beta_k^2} \mathbf{E}_{\tilde{\delta}_k} [\|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0\|^2] \\
&\quad - \eta_k \beta_k \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\mathbf{K}}^2 \\
&\quad + \frac{1}{2} \eta_k^2 \beta_k^2 \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\mathbf{K}}^2 + \frac{1}{2 \beta_k^2} \mathbf{E}_{\tilde{\delta}_k} [\|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0\|^2] \\
&\quad + \frac{1}{2} \eta_k \|\mathbf{g}_k - \mathbf{g}_k^0\|^2 + \frac{1}{2} \eta_k \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\mathbf{K}}^2 \\
&\quad + \frac{1}{2} \eta_k^2 \|\mathbf{g}_k - \mathbf{g}_k^0\|^2 + \frac{1}{2} \eta_k^2 \beta_k^2 \|\mathbf{L} \mathbf{x}_k\|^2 \\
&\quad + \frac{1}{2} \eta_k^2 \mathbf{E}_{\tilde{\delta}_k} [\|\mathbf{g}_k^\mu - \mathbf{g}_k^0\|^2] + \frac{1}{2 \beta_k^2} \mathbf{E}_{\tilde{\delta}_k} [\|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0\|^2] \\
&= \mathbf{x}_k^T (\mathbf{K} - \eta_k \alpha_k \mathbf{L}) \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right)
\end{aligned}$$

$$\begin{aligned}
&+ \frac{1}{2} (\eta_k + \eta_k^2) \|\mathbf{g}_k - \mathbf{g}_k^0\|^2 + \frac{1}{2} \eta_k^2 \mathbf{E}_{\tilde{\delta}_k} [\|\mathbf{g}_k^\mu - \mathbf{g}_k^0\|^2] \\
&+ \|\mathbf{x}_k\|_{\eta_k (\beta_k \mathbf{L} + \frac{1}{2} \mathbf{K}) + \eta_k^2 (\frac{1}{2} \alpha_k^2 - \alpha_k \beta_k + \beta_k^2) \mathbf{L}^2}^2 \\
&+ \left(\frac{1}{2 \eta_k \beta_k^2} + \frac{3}{2 \beta_k^2} \right) \mathbf{E}_{\tilde{\delta}_k} [\|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0\|^2] \\
&- \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\eta_k (\beta_k - \frac{1}{2} - \eta_k \beta_k^2) \mathbf{K}}^2 \\
&\leq \mathbf{x}_k^T \mathbf{K} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) - (1 + \omega_k) \eta_k \alpha_k \mathbf{x}_k^T \mathbf{L} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \\
&\quad + \omega_k \eta_k \alpha_k \mathbf{x}_k^T \mathbf{L} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \\
&\quad + \|\mathbf{x}_k\|_{\eta_k (\beta_k \mathbf{L} + \frac{1}{2} \mathbf{K}) + \eta_k^2 (\frac{1}{2} \alpha_k^2 - \alpha_k \beta_k + \beta_k^2) \mathbf{L}^2 + \frac{1}{2} \eta_k (1 + 3 \eta_k) \mathbf{L}_f^2 \mathbf{K}}^2 \\
&\quad + \frac{\eta_k}{2 \beta_k^2} (1 + 3 \eta_k) \mathbf{L}_f^2 \mathbf{E}_{\tilde{\delta}_k} [\|\bar{\mathbf{g}}_k^\mu\|^2] + n \sigma^2 \eta_k^2 \\
&\quad - \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\eta_k (\beta_k - \frac{1}{2} - \eta_k \beta_k^2) \mathbf{K}}^2 \tag{62}
\end{aligned}$$

where the first equality holds due to (5); the second equality holds since (25) in Lemma 3, \mathbf{x}_k and \mathbf{v}_k are independent of $\tilde{\delta}_k$, and (48a); the first inequality holds due to the Cauchy-Schwarz inequality, (25), $\rho(\mathbf{K}) = 1$, and the Jensen's inequality; and the last inequality holds due to (47), (49), and (55). For the third term on the right-hand side of (62), we have

$$\begin{aligned}
&\omega_k \eta_k \alpha_k \mathbf{x}_k^T \mathbf{L} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \\
&= \omega_k \eta_k \alpha_k \mathbf{x}_k^T \mathbf{L} \mathbf{K} \left(\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right) \\
&\leq \|\mathbf{x}_k\|_{\frac{1}{2} \omega_k \eta_k \alpha_k \mathbf{L}^2}^2 + \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\frac{1}{2} \omega_k \eta_k \alpha_k \mathbf{K}}^2. \tag{63}
\end{aligned}$$

Then, from (61)–(63), we have (60). \blacksquare

Lemma 9: Suppose Assumptions 2–5 hold. Then the following holds for Algorithm 1:

$$\begin{aligned}
\mathbf{E}_{\tilde{\delta}_k} [W_{4,k+1}] &\leq W_{4,k} - \frac{\eta_k}{4} \|\bar{\mathbf{g}}_k\|^2 + \|\mathbf{x}_k\|_{\frac{\eta_k}{2} \mathbf{L}_f^2 \mathbf{K}}^2 \\
&\quad - \frac{\eta_k}{4} \|\bar{\mathbf{g}}_k^0\|^2 + \frac{1}{2} \eta_k^2 \mathbf{L}_f \mathbf{E}_{\tilde{\delta}_k} [\|\bar{\mathbf{g}}_k^\mu\|^2] \tag{64}
\end{aligned}$$

where $W_{4,k} = n(f(\bar{\mathbf{x}}_k) - f^*) = \tilde{f}(\bar{\mathbf{x}}_k) - n f^*$.

Proof: We first note that $W_{4,k}$ is well defined due to $f^* > -\infty$ as assumed in Assumption 2.

From (47) and $\rho(\mathbf{H}) = 1$, it can be obtained that

$$\|\bar{\mathbf{g}}_k^0 - \bar{\mathbf{g}}_k\|^2 = \|\mathbf{H}(\mathbf{g}_k^0 - \mathbf{g}_k)\|^2 \leq \|\mathbf{g}_k^0 - \mathbf{g}_k\|^2 \leq \mathbf{L}_f^2 \|\mathbf{x}_k\|_{\mathbf{K}}^2. \tag{65}$$

From (48a), it can be calculated that

$$\mathbf{E}_{\tilde{\delta}_k} [\bar{\mathbf{g}}_k^\mu] = \mathbf{E}_{\tilde{\delta}_k} [\mathbf{H} \mathbf{g}_k^\mu] = \mathbf{H} \mathbf{E}_{\tilde{\delta}_k} [\mathbf{g}_k^\mu] = \bar{\mathbf{g}}_k. \tag{66}$$

We have

$$\begin{aligned}
\mathbf{E}_{\tilde{\delta}_k} [W_{4,k+1}] &= \mathbf{E}_{\tilde{\delta}_k} [\tilde{f}(\bar{\mathbf{x}}_{k+1}) - n f^*] \\
&= \mathbf{E}_{\tilde{\delta}_k} [\tilde{f}(\bar{\mathbf{x}}_k) - n f^* + \tilde{f}(\bar{\mathbf{x}}_{k+1}) - \tilde{f}(\bar{\mathbf{x}}_k)]
\end{aligned}$$

$$\begin{aligned}
 &\leq \mathbf{E}_{\tilde{\delta}_k}[\tilde{f}(\bar{\mathbf{x}}_k) - nf^* - \eta_k(\bar{\mathbf{g}}_k^u)^T \mathbf{g}_k^0 + \frac{1}{2}\eta_k^2 L_f \|\bar{\mathbf{g}}_k^u\|^2] \\
 &= \tilde{f}(\bar{\mathbf{x}}_k) - nf^* - \eta_k \bar{\mathbf{g}}_k^T \mathbf{g}_k^0 + \frac{1}{2}\eta_k^2 L_f \mathbf{E}_{\tilde{\delta}_k}[\|\bar{\mathbf{g}}_k^u\|^2] \\
 &= \tilde{f}(\bar{\mathbf{x}}_k) - nf^* - \eta_k \bar{\mathbf{g}}_k^T \bar{\mathbf{g}}_k^0 + \frac{1}{2}\eta_k^2 L_f \mathbf{E}_{\tilde{\delta}_k}[\|\bar{\mathbf{g}}_k^u\|^2] \\
 &= W_{4,k} - \frac{\eta_k}{2} \bar{\mathbf{g}}_k^T (\bar{\mathbf{g}}_k + \bar{\mathbf{g}}_k^0 - \bar{\mathbf{g}}_k) \\
 &\quad - \frac{\eta_k}{2} (\bar{\mathbf{g}}_k - \bar{\mathbf{g}}_k^0 + \bar{\mathbf{g}}_k^0)^T \bar{\mathbf{g}}_k^0 + \frac{1}{2}\eta_k^2 L_f \mathbf{E}_{\tilde{\delta}_k}[\|\bar{\mathbf{g}}_k^u\|^2] \\
 &\leq W_{4,k} - \frac{\eta_k}{4} \|\bar{\mathbf{g}}_k\|^2 + \frac{\eta_k}{4} \|\bar{\mathbf{g}}_k^0 - \bar{\mathbf{g}}_k\|^2 - \frac{\eta_k}{4} \|\bar{\mathbf{g}}_k^0\|^2 \\
 &\quad + \frac{\eta_k}{4} \|\bar{\mathbf{g}}_k^0 - \bar{\mathbf{g}}_k\|^2 + \frac{1}{2}\eta_k^2 L_f \mathbf{E}_{\tilde{\delta}_k}[\|\bar{\mathbf{g}}_k^u\|^2] \\
 &= W_{4,k} - \frac{\eta_k}{4} \|\bar{\mathbf{g}}_k\|^2 + \frac{\eta_k}{2} \|\bar{\mathbf{g}}_k^0 - \bar{\mathbf{g}}_k\|^2 \\
 &\quad - \frac{\eta_k}{4} \|\bar{\mathbf{g}}_k^0\|^2 + \frac{1}{2}\eta_k^2 L_f \mathbf{E}_{\tilde{\delta}_k}[\|\bar{\mathbf{g}}_k^u\|^2] \tag{67}
 \end{aligned}$$

where the first inequality holds since that \tilde{f} is smooth, (24a) and (54); the third equality holds since \mathbf{x}_k and \mathbf{v}_k are independent of $\tilde{\delta}_k$ and (66); the fourth equality holds due to $\bar{\mathbf{g}}_k^T \mathbf{g}_k^0 = \mathbf{g}_k^T \mathbf{H} \mathbf{g}_k^0 = \mathbf{g}_k^T \mathbf{H} \mathbf{H} \mathbf{g}_k^0 = \bar{\mathbf{g}}_k^T \bar{\mathbf{g}}_k^0$; and the second inequality holds due to the Cauchy-Schwarz inequality.

Then, from (65) and (67), we have (64). \blacksquare

B. Proof of Theorem 1

We denote the following notations:

$$c_0(\kappa_1, \kappa_2) = \max\{4\kappa_2\varepsilon_5, \varepsilon_6\}$$

$$c_1 = \frac{1}{\rho_2(L)} + 1$$

$$c_2(\kappa_1) = \min\left\{\frac{\varepsilon_1}{\varepsilon_2}, \frac{1}{5}\right\}$$

$$\kappa_3 = \frac{1}{\rho_2(L)} + \kappa_1 + 1$$

$$\kappa_4 = \frac{1}{\rho_2(L)} + \kappa_1 + \frac{3}{2}$$

$$\kappa_5 = \frac{\kappa_1 + 1}{2} + \frac{1}{2\rho_2(L)}$$

$$\kappa_6 = \min\left\{\frac{1}{2\rho(L)}, \frac{\kappa_1 - 1}{2\kappa_1}\right\}$$

$$\varepsilon_1 = (\kappa_1 - 1)\rho_2(L) - 1$$

$$\varepsilon_2 = \rho(L) + (2\kappa_1^2 + 1)\rho(L^2) + 1$$

$$\varepsilon_3 = \varepsilon_1\kappa_2 - \varepsilon_2\kappa_2^2$$

$$\varepsilon_4 = \frac{1}{2}(\kappa_2 - 5\kappa_2^2)$$

$$\varepsilon_5 = L_f + \frac{1}{\kappa_2\varepsilon_6}\kappa_3L_f^2 + \frac{2}{\varepsilon_6^2}\kappa_4L_f^2$$

$$\varepsilon_6 = \max\left\{\frac{1}{2}(2 + 3L_f^2), \kappa_3\right\}.$$

To prove Theorem 1, we need the following lemma:

Lemma 10: Suppose Assumptions 1–6 hold. Suppose $\alpha_k = \alpha = \kappa_1\beta$, $\beta_k = \beta \geq c_0(\kappa_1, \kappa_2)$, and $\eta_k = \eta = \kappa_2/\beta$, where $\kappa_1 > c_1$ and $\kappa_2 \in (0, c_2(\kappa_1))$. Then, for any $k \in \mathbb{N}_0$ the following holds for Algorithm 1:

$$\begin{aligned}
 \mathbf{E}_{\tilde{\delta}_k}[W_{k+1}] &\leq W_k - \|\mathbf{x}_k\|_{\varepsilon_3\mathbf{K}}^2 - \left\|\mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0\right\|_{\varepsilon_4\mathbf{K}}^2 \\
 &\quad - \frac{1}{4}\eta\|\bar{\mathbf{g}}_k^0\|^2 + (\varepsilon_5 + 3n)\sigma^2\eta^2 \tag{68a}
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{E}_{\tilde{\delta}_k}[\check{W}_{k+1}] &\leq \check{W}_k - \|\mathbf{x}_k\|_{\varepsilon_3\mathbf{K}}^2 - \left\|\mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0\right\|_{\varepsilon_4\mathbf{K}}^2 + 2\varepsilon_5\eta^2\|\bar{\mathbf{g}}_k^0\|^2 \\
 &\quad + 2L_f^2\varepsilon_5\eta^2\|\mathbf{x}_k\|_{\mathbf{K}}^2 + (\varepsilon_5 + 3n)\sigma^2\eta^2 \tag{68b}
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{E}_{\tilde{\delta}_k}[W_{4,k+1}] &\leq W_{4,k} - \frac{1}{4}\eta\|\bar{\mathbf{g}}_k^0\|^2 + \|\mathbf{x}_k\|_{\frac{1}{2}\eta L_f^2\mathbf{K}}^2 + L_f\sigma^2\eta^2 \tag{68c}
 \end{aligned}$$

where $W_k = \sum_{i=1}^4 W_{i,k}$ and $\check{W}_k = \sum_{i=1}^3 W_{i,k}$.

Proof: i) Noting that $\alpha_k = \alpha = \kappa_1\beta$, $\beta_k = \beta$, $\eta_k = \eta$, and $\omega_k = \frac{1}{\beta_k} - \frac{1}{\beta_{k+1}} = 0$, from (46), (51), (60), and (64), we have

$$\begin{aligned}
 \mathbf{E}_{\tilde{\delta}_k}[W_{k+1}] &\leq W_k + \left\|\mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0\right\|_{\frac{3}{2}\eta^2\beta^2\mathbf{K}}^2 + 2n\sigma^2\eta^2 \\
 &\quad - \|\mathbf{x}_k\|_{\eta\alpha L - \frac{1}{2}\eta\mathbf{K} - \frac{3}{2}\eta^2\alpha^2L^2 - \frac{1}{2}\eta(1+5\eta)L_f^2\mathbf{K}}^2 \\
 &\quad + \|\mathbf{x}_k\|_{\eta^2\beta^2(L+\kappa_1L^2)}^2 + \frac{1}{2}\eta\left(\frac{1}{\rho_2(L)} + \kappa_1\right)\left\|\mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0\right\|_{\mathbf{K}}^2 \\
 &\quad + \frac{\eta}{\beta^2}\left(\eta + \frac{1}{2}\right)\left(\frac{1}{\rho_2(L)} + \kappa_1\right)L_f^2\mathbf{E}_{\tilde{\delta}_k}[\|\bar{\mathbf{g}}_k^u\|^2] \\
 &\quad + \|\mathbf{x}_k\|_{\eta(\beta L + \frac{1}{2}\mathbf{K}) + \eta^2(\frac{1}{2}\alpha^2 - \alpha\beta + \beta^2)L^2 + \frac{1}{2}\eta(1+3\eta)L_f^2\mathbf{K}}^2 \\
 &\quad + \frac{\eta}{2\beta^2}(1 + 3\eta)L_f^2\mathbf{E}_{\tilde{\delta}_k}[\|\bar{\mathbf{g}}_k^u\|^2] + n\sigma^2\eta^2 \\
 &\quad - \left\|\mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0\right\|_{\eta(\beta - \frac{1}{2} - \eta\beta^2)\mathbf{K}}^2 - \frac{1}{4}\eta\|\bar{\mathbf{g}}_k\|^2 \\
 &\quad + \|\mathbf{x}_k\|_{\frac{1}{2}\eta L_f^2\mathbf{K}}^2 - \frac{1}{4}\eta\|\bar{\mathbf{g}}_k^0\|^2 + \frac{1}{2}\eta^2 L_f \mathbf{E}_{\tilde{\delta}_k}[\|\bar{\mathbf{g}}_k^u\|^2]. \tag{69}
 \end{aligned}$$

Note that

$$\begin{aligned}
 \mathbf{E}_{\tilde{\delta}_k}[\|\bar{\mathbf{g}}_k^u\|^2] &= \mathbf{E}_{\tilde{\delta}_k}[\|\bar{\mathbf{g}}_k^u - \bar{\mathbf{g}}_k + \bar{\mathbf{g}}_k\|^2] \\
 &\leq 2\mathbf{E}_{\tilde{\delta}_k}[\|\bar{\mathbf{g}}_k^u - \bar{\mathbf{g}}_k\|^2] + 2\|\bar{\mathbf{g}}_k\|^2 \\
 &= 2n\mathbf{E}_{\tilde{\delta}_k}\left[\left\|\frac{1}{n}\sum_{i=1}^n (g_{i,k}^u - g_{i,k})\right\|^2\right] + 2\|\bar{\mathbf{g}}_k\|^2 \\
 &= \frac{2}{n}\mathbf{E}_{\tilde{\delta}_k}\left[\left\|\sum_{i=1}^n (g_{i,k}^u - g_{i,k})\right\|^2\right] + 2\|\bar{\mathbf{g}}_k\|^2 \\
 &= \frac{2}{n}\sum_{i=1}^n \mathbf{E}_{\tilde{\delta}_k}[\|g_{i,k}^u - g_{i,k}\|^2] + 2\|\bar{\mathbf{g}}_k\|^2 \\
 &\leq 2\sigma^2 + 2\|\bar{\mathbf{g}}_k\|^2 \tag{70}
 \end{aligned}$$

where the first inequality holds due to the Cauchy-Schwarz inequality; the last equality holds since $\{g_{i,k}^u, i \in [n]\}$ are

independent of each other as assumed in Assumption 4, \mathbf{x}_k and \mathbf{v}_k are independent of $\tilde{\mathcal{F}}_k$, and $\mathbf{E}_{\tilde{\mathcal{F}}_k}[g_{i,k}^u] = g_{i,k}$ as assumed in Assumption 5; and the last inequality holds due to (48b).

From (69), (70), and $\alpha = \kappa_1\beta$, we have

$$\begin{aligned} \mathbf{E}_{\tilde{\mathcal{F}}_k}[W_{k+1}] &\leq W_k - \|\mathbf{x}_k\|_{\eta\mathbf{M}_1 - \eta^2\mathbf{M}_2}^2 - \left\| \mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0 \right\|_{b_{1,k}\mathbf{K}}^2 \\ &\quad - b_{2,k}\eta\|\tilde{\mathbf{g}}_k\|^2 - \frac{1}{4}\eta\|\tilde{\mathbf{g}}_k^0\|^2 + b_{3,k}\sigma^2\eta^2 + 3n\sigma^2\eta^2 \end{aligned} \quad (71)$$

where

$$\begin{aligned} \mathbf{M}_1 &= (\alpha - \beta)\mathbf{L} - \frac{1}{2}(2 + 3L_f^2)\mathbf{K} \\ \mathbf{M}_2 &= \beta^2\mathbf{L} + (2\alpha^2 + \beta^2)\mathbf{L}^2 + 4L_f^2\mathbf{K} \\ b_{1,k} &= \frac{1}{2}(2\beta - \kappa_3)\eta - \frac{5}{2}\beta^2\eta^2 \\ b_{2,k} &= \frac{1}{4} - b_{3,k}\eta \\ b_{3,k} &= L_f + \frac{1}{\beta^2\eta}\kappa_3L_f^2 + \frac{2}{\beta^2}\kappa_4L_f^2. \end{aligned}$$

From (26), $\alpha = \kappa_1\beta$, $\kappa_1 > c_1 > 1$, $\eta = \kappa_2/\beta$, and $\beta \geq c_0(\kappa_1, \kappa_2) \geq \varepsilon_6 \geq (2 + 3L_f^2)/2$, we have

$$\eta\mathbf{M}_1 \geq \varepsilon_1\kappa_2\mathbf{K}. \quad (72)$$

From (26), $\alpha = \kappa_1\beta$, and $\beta \geq \frac{1}{2}(2 + 3L_f^2) > 2L_f$, one can have

$$\eta^2\mathbf{M}_2 \leq \varepsilon_2\kappa_2^2\mathbf{K}. \quad (73)$$

From $\beta \geq \kappa_3$, it can be obtained that

$$b_{1,k} \geq \varepsilon_4. \quad (74)$$

From $\kappa_1 > c_1 = 1/\rho_2(L) + 1$, it can be derived that

$$\varepsilon_1 > 0. \quad (75)$$

From (75) and $\kappa_2 \in (0, \min\{\frac{\varepsilon_1}{\varepsilon_2}, \frac{1}{5}\})$, we have

$$\varepsilon_3 > 0 \quad (76a)$$

$$\varepsilon_4 > 0. \quad (76b)$$

From (76a), (76b), and $\beta \geq 4\kappa_2\varepsilon_5$, we have

$$b_{3,k} = L_f + \frac{1}{\beta^2\eta}\kappa_3L_f^2 + \frac{2}{\beta^2}\kappa_4L_f^2 \leq \varepsilon_5 \quad (77a)$$

$$b_{2,k} = \frac{1}{4} - b_{3,k}\eta \geq \frac{1}{4} - \frac{\kappa_2}{\beta}\varepsilon_5 \geq 0. \quad (77b)$$

From (71)–(74), (77a), and (77b), we have (68a).

ii) Similarly, the way to get (68a), we know that

$$\begin{aligned} \mathbf{E}_{\tilde{\mathcal{F}}_k}[\check{W}_{k+1}] &\leq \check{W}_k - \|\mathbf{x}_k\|_{\varepsilon_3\mathbf{K}}^2 - \left\| \mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0 \right\|_{\varepsilon_4\mathbf{K}}^2 \\ &\quad + \varepsilon_5\eta^2\|\tilde{\mathbf{g}}_k\|^2 + (\varepsilon_5 + 3n)\sigma^2\eta^2. \end{aligned} \quad (78)$$

We have

$$\begin{aligned} \|\tilde{\mathbf{g}}_k\|^2 &= \|\tilde{\mathbf{g}}_k - \tilde{\mathbf{g}}_k^0 + \tilde{\mathbf{g}}_k^0\|^2 \\ &\leq 2\|\tilde{\mathbf{g}}_k - \tilde{\mathbf{g}}_k^0\|^2 + 2\|\tilde{\mathbf{g}}_k^0\|^2 \leq 2L_f^2\|\mathbf{x}_k\|_{\mathbf{K}}^2 + 2\|\tilde{\mathbf{g}}_k^0\|^2 \end{aligned} \quad (79)$$

where the last inequality holds due to (65).

From (78) and (79), we have (68b).

iii) From (64) and (70), we have

$$\begin{aligned} \mathbf{E}_{\tilde{\mathcal{F}}_k}[W_{4,k+1}] &\leq W_{4,k} - \frac{1}{4}\eta\|\tilde{\mathbf{g}}_k\|^2 + \|\mathbf{x}_k\|_{\frac{1}{2}\eta L_f^2\mathbf{K}}^2 \\ &\quad - \frac{1}{4}\eta\|\tilde{\mathbf{g}}_k^0\|^2 + \eta^2L_f(\sigma^2 + \|\tilde{\mathbf{g}}_k\|^2). \end{aligned} \quad (80)$$

From $\eta = \kappa_2/\beta$ and $\beta \geq 4\kappa_2\varepsilon_5 > 4\kappa_2L_f$, we have

$$\eta L_f < \frac{1}{4}. \quad (81)$$

From (80) and (81), we have (68c). ■

Now we are ready to prove Theorem 1.

Denote

$$\hat{V}_k = \|\mathbf{x}_k\|_{\mathbf{K}}^2 + \left\| \mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0 \right\|_{\mathbf{K}}^2 + n(f(\bar{x}_k) - f^*).$$

We have

$$\begin{aligned} W_k &= \frac{1}{2}\|\mathbf{x}_k\|_{\mathbf{K}}^2 + \frac{1}{2}\left\| \mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0 \right\|_{\mathbf{Q} + \kappa_1\mathbf{K}}^2 \\ &\quad + \mathbf{x}_k^T\mathbf{K}\left(\mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0\right) + n(f(\bar{x}_k) - f^*) \\ &\geq \frac{1}{2}\|\mathbf{x}_k\|_{\mathbf{K}}^2 + \frac{1}{2}\left(\frac{1}{\rho(L)} + \kappa_1\right)\left\| \mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0 \right\|_{\mathbf{K}}^2 \\ &\quad - \frac{1}{2\kappa_1}\|\mathbf{x}_k\|_{\mathbf{K}}^2 - \frac{\kappa_1}{2}\left\| \mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0 \right\|_{\mathbf{K}}^2 + n(f(\bar{x}_k) - f^*) \\ &\geq \kappa_6\left(\|\mathbf{x}_k\|_{\mathbf{K}}^2 + \left\| \mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0 \right\|_{\mathbf{K}}^2\right) + n(f(\bar{x}_k) - f^*) \\ &\geq \kappa_6\hat{V}_k \geq 0 \end{aligned} \quad (82)$$

where the first inequality holds due to (28) and the Cauchy-Schwarz inequality; and the last inequality holds due to $0 < \kappa_6 < 0.5$. Similarly, we have

$$W_k \leq \kappa_5\hat{V}_k. \quad (84)$$

From (68a) and (76b), we have

$$\mathbf{E}_{\tilde{\mathcal{F}}_k}[W_{k+1}] \leq W_k - \varepsilon_3\|\mathbf{x}_k\|_{\mathbf{K}}^2 - \frac{\kappa_2}{4\beta}\|\tilde{\mathbf{g}}_k^0\|^2 + \frac{(\varepsilon_5 + 3n)\kappa_2^2\sigma^2}{\beta^2}. \quad (85)$$

Then, taking expectation in \mathcal{F}_T and summing (85) over $k \in [0, T]$ yield

$$\begin{aligned} \mathbf{E}[W_{T+1}] &+ \sum_{k=0}^T \mathbf{E}\left[\varepsilon_3\|\mathbf{x}_k\|_{\mathbf{K}}^2 + \frac{\kappa_2}{4\beta}\|\tilde{\mathbf{g}}_k^0\|^2\right] \\ &\leq W_0 + \frac{(T+1)(\varepsilon_5 + 3n)\kappa_2^2\sigma^2}{\beta^2}. \end{aligned} \quad (86)$$

From (86), (83), and (76a), we have

$$\begin{aligned} \frac{1}{T+1} \sum_{k=0}^T \mathbf{E}\left[\frac{1}{n} \sum_{i=1}^n \|x_{i,k} - \bar{x}_k\|^2\right] &= \frac{1}{n(T+1)} \sum_{k=0}^T \mathbf{E}[\|\mathbf{x}_k\|_{\mathbf{K}}^2] \\ &\leq \frac{W_0}{n\varepsilon_3(T+1)} + \frac{(\varepsilon_5 + 3n)\kappa_2^2\sigma^2}{n\varepsilon_3\beta^2}. \end{aligned} \quad (87)$$

Noting that $W_0 = \mathcal{O}(n)$, from (87), we have (11a).

Taking expectation in \mathcal{F}_T and summing (68c) over $k \in [0, T]$

yield

$$\begin{aligned} \frac{1}{4}n \sum_{k=0}^T \mathbf{E}[\|\nabla f(\bar{x}_k)\|^2] &= \frac{1}{4} \sum_{k=0}^T \mathbf{E}[\|\bar{\mathbf{g}}_k^0\|^2] \\ &\leq \frac{W_{4,0}}{\eta} + \frac{1}{2}L_f^2 \sum_{k=0}^T \mathbf{E}[\|\mathbf{x}_k\|_K^2] + (T+1)L_f\sigma^2\eta. \end{aligned} \quad (88)$$

From (88), $\eta = \kappa_2/\beta = \sqrt{n}/\sqrt{T}$, and (87), we have

$$\begin{aligned} \frac{1}{T} \sum_{k=0}^{T-1} \mathbf{E}[\|\nabla f(\bar{x}_k)\|^2] &\leq \frac{4\beta}{\kappa_2 T} (f(\bar{x}_0) - f^*) + \frac{4L_f\sigma^2\kappa_2}{n\beta} \\ &\quad + O\left(\frac{1}{T}\right) + O\left(\frac{1}{\beta^2}\right) \end{aligned}$$

which gives (11b).

Taking expectation in \mathcal{F}_T and summing (68c) over $k \in [0, T]$ yield

$$\begin{aligned} n(\mathbf{E}[f(\bar{x}_{T+1})] - f^*) &= \mathbf{E}[W_{4,T+1}] \\ &\leq W_{4,0} + \frac{1}{2}\eta L_f^2 \sum_{k=0}^T \mathbf{E}[\|\mathbf{x}_k\|_K^2] + L_f\sigma^2\eta^2(T+1). \end{aligned} \quad (89)$$

From (86), (89), and $\eta = \kappa_2/\beta$, we have (11c). \blacksquare

C. Proof of Theorem 2

In addition to the notations defined in Appendix B, we also denote

$$\varepsilon_7 = \frac{1}{\kappa_5} \min\left\{\varepsilon_3, \varepsilon_4, \frac{\nu}{2(T+1)^\theta}\right\}.$$

From the conditions in Theorem 2, we know that all conditions needed in Lemma 10 are satisfied, so (68a)–(68c) still hold.

From Assumptions 2 and 7 as well as (13), we have that

$$\|\bar{\mathbf{g}}_k^0\|^2 = n\|\nabla f(\bar{x}_k)\|^2 \geq 2\nu n(f(\bar{x}_k) - f^*) = 2\nu W_{4,k}. \quad (90)$$

From (83), we have

$$\|\mathbf{x}_k\|_K^2 + W_{4,k} \leq \hat{v}_k \leq \frac{W_k}{\kappa_6}. \quad (91)$$

From (68a), (90), (83), (84), and (14), we have

$$\begin{aligned} \mathbf{E}_{\bar{\mathcal{F}}_k}[W_{k+1}] &\leq W_k - \varepsilon_3\|\mathbf{x}_k\|_K^2 - \varepsilon_4\left\|\mathbf{v}_k + \frac{1}{\beta_k}\mathbf{g}_k^0\right\|_K^2 \\ &\quad - \frac{1}{2}\eta\nu W_{4,k} + (\varepsilon_5 + 3n)\sigma^2\eta^2 \\ &\leq W_k - \frac{1}{\kappa_5} \min\left\{\varepsilon_3, \varepsilon_4, \frac{\nu\eta}{2}\right\}W_k + (\varepsilon_5 + 3n)\sigma^2\eta^2. \end{aligned} \quad (92)$$

From (92) and (14), we have

$$\mathbf{E}_{\bar{\mathcal{F}}_k}[W_{k+1}] \leq W_k - \varepsilon_7 W_k + \frac{(\varepsilon_5 + 3n)\sigma^2}{(T+1)^{2\theta}}, \quad \forall k \leq T. \quad (93)$$

From $\kappa_1 > 1$ and $\rho_2(L) > 0$, we have $\kappa_5 = \frac{\kappa_1+1}{2} + \frac{1}{2\rho_2(L)} > 1$. From $0 < \kappa_2 < c_2(\kappa_1) = \min\left\{\frac{\varepsilon_1}{\varepsilon_2}, \frac{1}{5}\right\} < 1/5$, we have $\varepsilon_4 = (\kappa_2 - 5\kappa_2^2)/2 \leq \max_{\kappa_2 \in (0, 1/5)} (\kappa_2 - 5\kappa_2^2)/2 = 1/40$. Thus,

$$0 < \varepsilon_7 = \frac{1}{\kappa_5} \min\left\{\varepsilon_3, \varepsilon_4, \frac{\nu}{2(T+1)^\theta}\right\} \leq \frac{\varepsilon_4}{\kappa_5} \leq \frac{1}{40}. \quad (94)$$

Then, from (93), (83), and (94), we have

$$\begin{aligned} \mathbf{E}[W_{k+1}] &\leq (1 - \varepsilon_7)^{k+1} W_0 + \frac{(\varepsilon_5 + 3n)\sigma^2}{(T+1)^{2\theta}} \sum_{l=0}^k (1 - \varepsilon_7)^l \\ &\leq (1 - \varepsilon_7)^{k+1} W_0 + \frac{(\varepsilon_5 + 3n)\sigma^2}{\varepsilon_7(T+1)^{2\theta}}, \quad \forall k \leq T. \end{aligned} \quad (95)$$

Then, noting that $\varepsilon_7 = O(1/(T+1)^\theta)$ and $\theta \in (0, 1)$, from (95), (29), and (91), we have

$$\mathbf{E}[\|\mathbf{x}_k\|_K^2 + W_{4,k}] = O\left(\frac{n}{T^\theta}\right), \quad \forall k \leq T. \quad (96)$$

Thus, there exists a constant $c_f > 0$ such that

$$\mathbf{E}[\|\mathbf{x}_k\|_K^2 + W_{4,k}] \leq nc_f, \quad \forall k \leq T. \quad (97)$$

From (82) and (84), we have

$$0 \leq 2\kappa_6(W_{1,k} + W_{2,k}) \leq \check{W}_k \leq 2\kappa_5(W_{1,k} + W_{2,k}). \quad (98)$$

From (24b), we have

$$\|\bar{\mathbf{g}}_k^0\|^2 = n\|\nabla f(\bar{x}_k)\|^2 \leq 2L_f n(f(\bar{x}_k) - f^*) = 2L_f W_{4,k}. \quad (99)$$

Denote $\check{z}_k = \mathbf{E}[\check{W}_k]$. From (68b) and (97)–(99), we have

$$\check{z}_{k+1} \leq (1 - a_1)\check{z}_k + a_2\eta^2, \quad \forall k \leq T \quad (100)$$

where $a_1 = \min\{\varepsilon_3, \varepsilon_4\}/\kappa_5$ and $a_2 = n(4\varepsilon_5 L_f c_f + 2\varepsilon_5 L_f^2 c_f + (\varepsilon_5 + 3)\sigma^2)$.

From (94), we have

$$a_1 \leq \frac{\varepsilon_4}{\kappa_5} \leq \frac{1}{40}. \quad (101)$$

From (100) and (101), we have

$$\check{z}_{k+1} \leq (1 - a_1)^{k+1} \check{z}_0 + \frac{a_2\eta^2}{a_1}, \quad \forall k \leq T$$

which yields (15a).

From (68c) and (90), we have

$$\begin{aligned} \mathbf{E}_{\bar{\mathcal{F}}_k}[W_{4,k+1}] &\leq \left(1 - \frac{\nu\eta}{2}\right)W_{4,k} + \frac{1}{2}\eta L_f^2 \|\mathbf{x}_k\|_K^2 + L_f\sigma^2\eta^2 \\ &\leq \left(1 - \frac{\nu\eta}{2}\right)^{k+1} W_{4,0} + \frac{1}{\nu}(L_f^2 \|\mathbf{x}_k\|_K^2 + 2L_f\sigma^2\eta). \end{aligned} \quad (102)$$

Noting $\eta = 1/(T+1)^\theta$, from (102), (29), and (15a), we have (15b). \blacksquare

D. Proof of Theorem 3

In addition to the notations defined in Appendix B, we also give the following notations:

$$\tilde{c}_0(\kappa_1, \kappa_2) = \max\left\{4\varepsilon_{11}, \varepsilon_6, \frac{\varepsilon_{10}}{\varepsilon_4}\right\}$$

$$\hat{c}_2(\kappa_1) = \min\left\{\frac{\varepsilon_1}{\varepsilon_2}, \frac{\varepsilon_8}{\varepsilon_9}, \frac{1}{5}\right\}$$

$$\hat{c}_3(\kappa_0, \kappa_1, \kappa_2) = \max\left\{\frac{\tilde{c}_0(\kappa_1, \kappa_2)}{\kappa_0}, \frac{8L_f\kappa_3}{\nu\kappa_2}, \frac{16L_f(\kappa_3 - 1)}{\nu\kappa_0\kappa_2}\right\}$$

$$\tilde{\sigma}^2 = 2L_f f^* - 2L_f \frac{1}{n} \sum_{i=1}^n f_i^*$$

$$\begin{aligned}
\varepsilon_8 &= \kappa_1 \rho_2(L) - 1 \\
\varepsilon_9 &= \frac{1}{2}(3\kappa_1 + 2)\kappa_1 \rho(L^2) + \rho(L) + 1 \\
\varepsilon_{10} &= \kappa_2(\kappa_3 - 1) + \kappa_1 \kappa_2 + \kappa_3 - 1 + 3\kappa_2^2 \\
\varepsilon_{11} &= \kappa_2 L_f + (2\kappa_3 - 1 + \kappa_2(10\kappa_3 - 4))L_f^2 \\
\varepsilon_{12} &= 3 + L_f + \frac{\kappa_3 L_f^2}{\kappa_0 \kappa_2 t_1} + \frac{2\kappa_4 L_f^2}{\kappa_0^2 t_1^2} + \frac{2 + 2\kappa_3 L_f^2}{\kappa_0 t_1^2} \\
&\quad + \frac{(\kappa_3 - 1)L_f^2}{\kappa_0^2 \kappa_2 t_1^3} + \frac{(\kappa_3 - 1)L_f^2}{\kappa_0^2 t_1^4} \left(\frac{2}{\kappa_0} + 2 \right) \\
\varepsilon_{13} &= \frac{\kappa_0 \kappa_3}{\kappa_2^2} + \frac{\kappa_3 - 1}{\kappa_2^2 t_1^2} \\
\varepsilon_{14} &= \varepsilon_{12} \sigma^2 + \varepsilon_{13} \tilde{\sigma}^2 \\
\varepsilon_{15} &= \frac{1}{\kappa_5} \min \left\{ \frac{\varepsilon_3 \kappa_0 t_1}{\kappa_2}, \frac{\varepsilon_4 \kappa_0 t_1}{2\kappa_2}, \frac{\nu}{8} \right\} \\
\varepsilon_{16} &= \frac{4L_f \sigma^2 \kappa_2^2}{\kappa_0^2 \left(\frac{\nu \kappa_2}{2\kappa_0} - 1 \right)}.
\end{aligned}$$

To prove Theorem 3, we need the following lemma.

Lemma 11: Suppose Assumptions 1–6 hold. Let $\alpha_k = \kappa_1 \beta_k$, $\beta_k = \kappa_0(k + t_1)$, and $\eta_k = \kappa_2/\beta_k$, where $\kappa_0 \geq \tilde{c}_0(\kappa_1, \kappa_2)/t_1$, $\kappa_1 > c_1$, $\kappa_2 \in (0, \hat{c}_2(\kappa_1))$, and $t_1 \geq 1$. Then, for any $k \in \mathbb{N}_0$ the following holds for Algorithm 1:

$$\begin{aligned}
\mathbf{E}_{\tilde{\delta}_k}[W_{k+1}] &\leq W_k - \varepsilon_3 \|\mathbf{x}_k\|_K^2 - \frac{1}{2} \varepsilon_4 \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_K^2 - \frac{1}{4} \eta_k \|\bar{\mathbf{g}}_k^0\|^2 \\
&\quad + 2L_f b_{8,k} \eta_k^2 W_{4,k} + n \varepsilon_{14} \eta_k^2
\end{aligned} \tag{103a}$$

$$\begin{aligned}
\mathbf{E}_{\tilde{\delta}_k}[\check{W}_{k+1}] &\leq \check{W}_k - \varepsilon_3 \|\mathbf{x}_k\|_K^2 - \frac{1}{2} \varepsilon_4 \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_K^2 + n \varepsilon_{14} \eta_k^2 \\
&\quad + 2\varepsilon_{12} L_f^2 \eta_k^2 \|\mathbf{x}_k\|_K^2 + 2(2\varepsilon_{12} + \varepsilon_{13}) L_f \eta_k^2 W_{4,k}
\end{aligned} \tag{103b}$$

$$\mathbf{E}_{\tilde{\delta}_k}[W_{4,k+1}] \leq W_{4,k} - \frac{\eta_k}{4} \|\bar{\mathbf{g}}_k^0\|^2 + \|\mathbf{x}_k\|_{\frac{1}{2}L_f^2 \eta_k \mathbf{K}}^2 + \eta_k^2 L_f \sigma^2 \tag{103c}$$

where $b_{8,k} = \kappa_3 \frac{\omega_k}{\eta_k^2} + (\kappa_3 - 1) \frac{\omega_k^2}{\eta_k^2}$.

Proof: i) We have

$$\begin{aligned}
\|\mathbf{g}_k^0\|^2 &= \sum_{i=1}^n \|\nabla f_i(\bar{\mathbf{x}}_k)\|^2 \leq \sum_{i=1}^n 2L_f (f_i(\bar{\mathbf{x}}_k) - f_i^*) \\
&= 2L_f n(f(\bar{\mathbf{x}}_k) - f^*) + n\tilde{\sigma}^2
\end{aligned} \tag{104}$$

where the inequality holds due to (24b).

From the Cauchy-Schwarz inequality, (55), and (104), we have

$$\begin{aligned}
\|\mathbf{g}_{k+1}^0\|^2 &= \|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0 + \mathbf{g}_k^0\|^2 \leq 2(\|\mathbf{g}_{k+1}^0 - \mathbf{g}_k^0\|^2 + \|\mathbf{g}_k^0\|^2) \\
&\leq 2(\eta_k^2 L_f^2 \|\bar{\mathbf{g}}_k^0\|^2 + 2L_f W_{4,k} + n\tilde{\sigma}^2).
\end{aligned} \tag{105}$$

From (46), (51), (60), (64), (70), (105), $\alpha_k = \kappa_1 \beta_k$, and $\eta_k = \kappa_2/\beta_k$, we have

$$\begin{aligned}
\mathbf{E}_{\tilde{\delta}_k}[W_{k+1}] &\leq W_k - \|\mathbf{x}_k\|_{\eta_k \mathbf{M}_{3,k} - \eta_k^2 \mathbf{M}_{4,k} - \frac{1}{2} \kappa_1 \kappa_2 \omega_k + \eta_k \omega_k \mathbf{M}_{5,k} - \eta_k^2 \omega_k \mathbf{M}_{6,k}}^2 \\
&\quad - \left\| \mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0 \right\|_{\beta_{4,k}^0 \mathbf{K}}^2 - \eta_k b_{5,k} \|\bar{\mathbf{g}}_k\|^2 - \frac{1}{4} \eta_k \|\bar{\mathbf{g}}_k^0\|^2 \\
&\quad + \eta_k^2 (b_{6,k} + b_{7,k} n) \sigma^2 + \eta_k^2 b_{8,k} (2L_f W_{4,k} + n\tilde{\sigma}^2)
\end{aligned} \tag{106}$$

where

$$\begin{aligned}
\mathbf{M}_{3,k} &= (\alpha_k - \beta_k) \mathbf{L} - \frac{1}{2} (2 + 3L_f^2) \mathbf{K} \\
\mathbf{M}_{4,k} &= \beta_k^2 \mathbf{L} + (2\alpha_k^2 + \beta_k^2) \mathbf{L}^2 + 4L_f^2 \mathbf{K} \\
\mathbf{M}_{5,k} &= \alpha_k \mathbf{L} - \frac{1}{2} (1 + L_f^2) \mathbf{K} \\
\mathbf{M}_{6,k} &= \frac{3}{2} \alpha_k^2 \mathbf{L}^2 + \beta_k^2 (\mathbf{L} + \kappa_1 \mathbf{L}^2) + \frac{5}{2} L_f^2 \mathbf{K} \\
b_{4,k}^0 &= \frac{1}{2} \eta_k (2\beta_k - \kappa_3) - \frac{5}{2} \eta_k^2 \beta_k^2 - \frac{1}{2} \omega_k \eta_k (\kappa_3 - 1) \\
&\quad - \frac{1}{2} \omega_k (\eta_k \alpha_k + \kappa_3 - 1 + 3\eta_k^2 \beta_k^2) \\
b_{5,k} &= \frac{1}{4} - \eta_k b_{6,k} \\
b_{6,k} &= L_f + \frac{1}{\beta_k^2 \eta_k} \kappa_3 L_f^2 + \frac{2}{\beta_k^2} \kappa_4 L_f^2 + 2\kappa_3 L_f^2 \omega_k \\
&\quad + \omega_k \left(\frac{1}{\beta_k^2 \eta_k} + \frac{2}{\beta_k^2} + 2\omega_k \right) (\kappa_3 - 1) L_f^2 \\
b_{7,k} &= 3 + 2\omega_k.
\end{aligned}$$

From (26), $\alpha_k = \kappa_1 \beta_k$, $\kappa_1 > 1$, $\beta_k \geq \kappa_0 t_1 \geq \tilde{c}_0(\kappa_1, \kappa_2) \geq \varepsilon_6 \geq (2 + 3L_f^2)/2$, and $\eta_k = \kappa_2/\beta_k$, we have

$$\eta_k \mathbf{M}_{3,k} \geq \varepsilon_1 \kappa_2 \mathbf{K}. \tag{107}$$

From (26), $\alpha_k = \kappa_1 \beta_k$, $\beta_k \geq (2 + 3L_f^2)/2 > 2L_f$, and $\eta_k = \kappa_2/\beta_k$, we have

$$\eta_k^2 \mathbf{M}_{4,k} \leq \varepsilon_2 \kappa_2^2 \mathbf{K}. \tag{108}$$

From (26), $\alpha_k = \kappa_1 \beta_k$, $\beta_k \geq (2 + 3L_f^2)/2 > (1 + L_f^2)/2$, and $\eta_k = \kappa_2/\beta_k$, we have

$$\eta_k \mathbf{M}_{5,k} \geq \varepsilon_8 \kappa_2 \mathbf{K}. \tag{109}$$

From (26), $\alpha_k = \kappa_1 \beta_k$, $\beta_k > 2L_f > \sqrt{10} L_f/2$, and $\eta_k = \kappa_2/\beta_k$, we have

$$\eta_k^2 \mathbf{M}_{6,k} \leq \varepsilon_9 \kappa_2^2 \mathbf{K}. \tag{110}$$

From $\alpha_k = \kappa_1 \beta_k$, $\beta_k \geq \kappa_3$, and $\eta_k = \kappa_2/\beta_k$, we have

$$b_{4,k}^0 \geq b_{4,k} \tag{111}$$

where $b_{4,k} = \varepsilon_4 - \frac{1}{2} \omega_k \eta_k (\kappa_3 - 1) - \frac{1}{2} \omega_k (\kappa_1 \kappa_2 + \kappa_3 - 1 + 3\kappa_2^2)$.

From $\kappa_1 > c_1 = 1/\rho_2(L) + 1$, we have

$$\varepsilon_1 > 0 \quad \text{and} \quad \varepsilon_8 > 0. \tag{112}$$

From (112) and $\kappa_2 \in (0, \min\{\frac{\varepsilon_1}{\varepsilon_2}, \frac{\varepsilon_8}{\varepsilon_9}, \frac{1}{5}\})$, we have

$$\varepsilon_3 > 0 \tag{113a}$$

$$\varepsilon_8 \kappa_2 - \varepsilon_9 \kappa_2^2 > 0 \tag{113b}$$

$$\varepsilon_4 > 0. \quad (113c)$$

From $\beta_k = \kappa_0(k + t_1)$, we have

$$\begin{aligned} \omega_k &= \frac{1}{\beta_k} - \frac{1}{\beta_{k+1}} = \frac{1}{\kappa_0} \left(\frac{1}{k+t_1} - \frac{1}{k+t_1+1} \right) \\ &= \frac{1}{\kappa_0(k+t_1)(k+t_1+1)} \leq \frac{\kappa_0}{\beta_k^2}. \end{aligned} \quad (114)$$

From (113a)–(114), and $\kappa_0 \geq \max\{\frac{4\varepsilon_{11}}{t_1}, \frac{\varepsilon_{10}}{\varepsilon_4 t_1}\}$, we have

$$b_{4,k} \geq \varepsilon_4 - \frac{\varepsilon_{10}}{2\kappa_0 t_1^2} \geq \varepsilon_4 - \frac{\varepsilon_{10}}{2\kappa_0 t_1} \geq \frac{1}{2}\varepsilon_4 > 0 \quad (115a)$$

$$b_{5,k} \geq \frac{1}{4} - \frac{\varepsilon_{11}}{\kappa_0 t_1} \geq 0. \quad (115b)$$

From (114) and $\beta_k = \kappa_0(k + t_1) \geq \kappa_0 t_1$, we have

$$b_{6,k} + b_{7,k} \leq \varepsilon_{12} \quad (116a)$$

$$b_{8,k} \leq \varepsilon_{13}. \quad (116b)$$

From (106)–(111), (113a)–(113c), and (115a)–(116b), we have (103a).

ii) Similarly, the way to get (103a), we have

$$\begin{aligned} \mathbf{E}_{\tilde{\delta}_k}[\check{W}_{k+1}] &\leq \check{W}_k - \varepsilon_3 \|\mathbf{x}_k\|_{\mathbf{K}}^2 - \frac{1}{2}\varepsilon_4 \|\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0\|_{\mathbf{K}}^2 + \varepsilon_{12} \eta_k^2 \|\bar{\mathbf{g}}_k\|^2 \\ &\quad + 2L_f \varepsilon_{13} \eta_k^2 W_{4,k} + n\varepsilon_{14} \eta_k^2, \quad \forall k \in \mathbb{N}_0. \end{aligned} \quad (117)$$

From (117), (79), and (99), we have (103b).

iii) From (64) and (70), we have

$$\begin{aligned} \mathbf{E}_{\tilde{\delta}_k}[W_{4,k+1}] &\leq W_{4,k} - \frac{\eta_k}{4} \|\bar{\mathbf{g}}_k\|^2 + \|\mathbf{x}_k\|_{\frac{1}{2}L_f^2 \eta_k \mathbf{K}}^2 \\ &\quad - \frac{\eta_k}{4} \|\bar{\mathbf{g}}_k^0\|^2 + \eta_k^2 L_f (\sigma^2 + \|\bar{\mathbf{g}}_k\|^2). \end{aligned} \quad (118)$$

From $0 < \eta_k \leq \kappa_2 / (\kappa_0 t_1)$ and $\kappa_0 t_1 \geq \tilde{c}_0(\kappa_1, \kappa_2) \geq 4\varepsilon_{11} > 4\kappa_2 L_f$, we have

$$\eta_k^2 L_f < \frac{1}{4} \eta_k. \quad (119)$$

From (118) and (119), we have (103c). \blacksquare

Now we are ready to prove Theorem 3.

From $t_1 > \hat{c}_3(\kappa_0, \kappa_1, \kappa_2) \geq \tilde{c}_0(\kappa_1, \kappa_2) / \kappa_0$, we have

$$\kappa_0 > \frac{\tilde{c}_0(\kappa_1, \kappa_2)}{t_1}.$$

Thus, all conditions needed in Lemma 11 are satisfied, so (103a)–(103c) hold.

From (103a) and (90), we have

$$\begin{aligned} \mathbf{E}_{\tilde{\delta}_k}[W_{k+1}] &\leq W_k - \varepsilon_3 \|\mathbf{x}_k\|_{\mathbf{K}}^2 - \frac{1}{2}\varepsilon_4 \|\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0\|_{\mathbf{K}}^2 - \frac{\eta_k \nu}{2} W_{4,k} \\ &\quad + 2L_f b_{8,k} \eta_k^2 W_{4,k} + n\varepsilon_{14} \eta_k^2 \\ &= W_k - \varepsilon_3 \|\mathbf{x}_k\|_{\mathbf{K}}^2 - \frac{1}{2}\varepsilon_4 \|\mathbf{v}_k + \frac{1}{\beta_k} \mathbf{g}_k^0\|_{\mathbf{K}}^2 + n\varepsilon_{14} \eta_k^2 \\ &\quad - 2\left(\frac{1}{4} - \frac{1}{\nu} L_f b_{8,k} \eta_k\right) \nu \eta_k W_{4,k}, \quad \forall k \in \mathbb{N}_0. \end{aligned} \quad (120)$$

From $t_1 > \hat{c}_3(\kappa_0, \kappa_1, \kappa_2) \geq 8L_f \kappa_3 / (\nu \kappa_2)$, we have

$$\frac{1}{4} - \frac{L_f \kappa_3}{\nu \kappa_2 t_1} \geq \frac{1}{8}. \quad (121)$$

From (114), (121), and $\kappa_0 > \tilde{c}_0(\kappa_1, \kappa_2) / t_1 \geq 16L_f(\kappa_3 - 1) / (\nu \kappa_2 t_1)$, we have

$$\begin{aligned} \frac{1}{4} - \frac{1}{\nu} L_f b_{8,k} \eta_k &\geq \frac{1}{4} - \frac{L_f \kappa_0 \kappa_3}{\nu \kappa_2 \beta_k} - \frac{L_f \kappa_0^2 (\kappa_3 - 1)}{\nu \kappa_2 \beta_k^3} \\ &\geq \frac{1}{4} - \frac{L_f \kappa_3}{\nu \kappa_2 t_1} - \frac{L_f (\kappa_3 - 1)}{\nu \kappa_2 \kappa_0 t_1^2} \geq \frac{1}{8} - \frac{L_f (\kappa_3 - 1)}{\nu \kappa_2 \kappa_0 t_1} \geq \frac{1}{16}. \end{aligned} \quad (122)$$

From (120), (83), and (84), we have

$$\begin{aligned} \mathbf{E}_{\tilde{\delta}_k}[W_{k+1}] &\leq W_k - \frac{\eta_k}{\kappa_5} \min\left\{\frac{\varepsilon_3}{\eta_k}, \frac{\varepsilon_4}{2\eta_k}, \frac{\nu}{8}\right\} W_k + n\varepsilon_{14} \eta_k^2 \\ &\leq W_k - \varepsilon_{15} \eta_k W_k + n\varepsilon_{14} \eta_k^2, \quad \forall k \in \mathbb{N}_0. \end{aligned} \quad (123)$$

Denote $z_k = \mathbf{E}[W_k]$, $r_{1,k} = \varepsilon_{15} \eta_k$, and $r_{2,k} = n\varepsilon_{14} \eta_k^2$, then from (123), we have

$$z_{k+1} \leq (1 - r_{1,k}) z_k + r_{2,k}, \quad \forall k \in \mathbb{N}_0. \quad (124)$$

From (16), we have

$$r_{1,k} = \eta_k \varepsilon_{15} = \frac{a_3}{k + t_1} \quad (125a)$$

$$r_{2,k} = \eta_k^2 \varepsilon_{14} n \sigma^2 = \frac{a_4}{(k + t_1)^2} \quad (125b)$$

where $a_3 = \kappa_2 \varepsilon_{15} / \kappa_0$ and $a_4 = n \kappa_2^2 \varepsilon_{14} / \kappa_0^2$.

From (94), we have

$$r_{1,k} \leq \frac{\varepsilon_4}{2\kappa_5} \leq \frac{1}{80}. \quad (126)$$

Then, from (124)–(126) and (32), we have

$$z_k \leq \phi_1(k, t_1, a_3, a_4, z_0), \quad \forall k \in \mathbb{N}_+ \quad (127)$$

where the function ϕ_1 is defined in (33).

From $\kappa_0 \geq \hat{c}_0 \nu \kappa_2 / 4$, we have

$$\phi_1(k, t_1, a_3, a_4, 2, z_0) = \begin{cases} \mathcal{O}\left(\frac{n}{k}\right), & \text{if } a_3 > 1 \\ \mathcal{O}\left(\frac{n \ln(k-1)}{k}\right), & \text{if } a_3 = 1 \\ \mathcal{O}\left(\frac{n}{k^{a_3}}\right), & \text{if } a_3 < 1. \end{cases} \quad (128)$$

From (127), (128), and (91), we know that there exists a constant $c_f > 0$ such that

$$\mathbf{E}[\|\mathbf{x}_k\|_{\mathbf{K}}^2 + W_{4,k}] \leq n c_f. \quad (129)$$

From (103b), (129), (98), and (16), we have

$$\check{z}_{k+1} \leq (1 - a_5) \check{z}_k + \frac{a_6}{(t + t_1)^2} \quad (130)$$

where $a_5 = \min\{\varepsilon_3, \varepsilon_4/2\} / \kappa_5$ and $a_6 = n(2\varepsilon_{12} L_f^2 c_f + 2(2\varepsilon_{12} + \varepsilon_{13}) L_f c_f + \varepsilon_{14}) \kappa_2^2 / \kappa_0^2$.

From (94), we have

$$a_5 \leq \frac{\varepsilon_4}{2\kappa_5} \leq \frac{1}{80}. \quad (131)$$

From (113a) and (113c), we know that

$$a_5 > 0 \quad \text{and} \quad a_6 > 0. \quad (132)$$

From (130)–(132) and (34), we have

$$\check{z}_k \leq \phi_2(k, t_1, a_5, a_6, \check{z}_0) = \mathcal{O}\left(\frac{n}{k^2}\right), \forall k \in \mathbb{N}_+ \quad (133)$$

where the function ϕ_2 is defined in (35).

From (35), (98), and (133), we have

$$\mathbf{E}[\|\mathbf{x}_k\|_{\mathbf{K}}^2] \leq \frac{1}{\kappa_6} \check{z}_k \leq \frac{1}{\kappa_6} \phi_2(k, t_1, a_5, a_6, \check{z}_0) = \mathcal{O}\left(\frac{n}{k^2}\right). \quad (134)$$

From (134), we have (17a).

From (103c) and (90), we have

$$\mathbf{E}[W_{4,k+1}] \leq \left(1 - \frac{\nu}{2}\eta\kappa\right)\mathbf{E}[W_{4,k}] + \|\mathbf{x}_k\|_{\frac{\nu}{2}L_f^2\eta\kappa\mathbf{K}}^2 + L_f\sigma^2\eta_k^2. \quad (135)$$

From $\kappa_0 < \nu\kappa_2/4$, we have

$$\frac{\nu\kappa_2}{2\kappa_0} > 2. \quad (136)$$

Similarly, the way to prove (32), from (134)–(136), we have

$$\mathbf{E}[f(\bar{\mathbf{x}}_T) - f^*] \leq \frac{\varepsilon_{16}}{n(T+t_1)} + \mathcal{O}\left(\frac{1}{(T+t_1)^2}\right) \quad (137)$$

where ε_{16} is determined by the last terms in (33) and (135).

From $\kappa_0 \geq \hat{c}_0\nu\kappa_2/4$, we have

$$\begin{aligned} \varepsilon_{16} &= \frac{4L_f\sigma^2\kappa_2^2}{\kappa_0^2\left(\frac{\nu\kappa_2}{2\kappa_0} - 1\right)} \leq \frac{4L_f\sigma^2\kappa_2^2}{\kappa_0^2\left(\frac{\nu\kappa_2}{2\kappa_0} - \frac{\nu\kappa_2}{4\kappa_0}\right)} \\ &= \frac{16L_f\sigma^2\kappa_2^2}{\nu\kappa_0} \leq \frac{64L_f\sigma^2}{\hat{c}_0\nu^2}. \end{aligned} \quad (138)$$

From (137) and (138), we have (17b). ■

E. Proof of Theorem 4

In addition to the notations defined in Appendices B and C, we also denote the following notations:

$$\varepsilon = \frac{1}{\kappa_5} \min\left\{\frac{\varepsilon_3}{\eta}, \frac{\varepsilon_4}{\eta}, \frac{\nu}{2}\right\}$$

$$c_4 = \frac{W_0}{n\kappa_6}$$

$$c_5 = \frac{\varepsilon_5 + 3n}{n\varepsilon\kappa_6}.$$

From the conditions in Theorem 4, we know that (92) holds. Thus,

$$\mathbf{E}_{\check{\delta}_k}[W_{k+1}] \leq W_k - \eta\varepsilon W_k + (\varepsilon_5 + 3n)\sigma^2\eta^2. \quad (139)$$

Similarly, the way to get (94), we have

$$0 < \eta\varepsilon < 1. \quad (140)$$

From (139) and (140), we have

$$\begin{aligned} \mathbf{E}[W_{k+1}] &\leq (1 - \eta\varepsilon)\mathbf{E}[W_k] + (\varepsilon_5 + 3n)\sigma^2\eta^2 \\ &\leq (1 - \eta\varepsilon)^{k+1}W_0 + (\varepsilon_5 + 3n)\sigma^2\eta^2 \sum_{\tau=0}^k (1 - \eta\varepsilon)^\tau \\ &\leq (1 - \eta\varepsilon)^{k+1}W_0 + \frac{\eta(\varepsilon_5 + 3n)\sigma^2}{\varepsilon}. \end{aligned} \quad (141)$$

Hence, (141) and (91) give (19). ■

F. Proof of Theorem 5

In addition to the notations defined in Appendices C, B, and E, we also denote the following notations:

$$\check{c}_0(\kappa_1, \kappa_2) = \max\{4\kappa_2\varepsilon_5, \check{\varepsilon}_6\}$$

$$\check{\varepsilon}_6 = \max\{1 + 3L_f^2, \kappa_3\}$$

$$\check{c}_5 = \frac{3 + 5\eta}{\varepsilon\kappa_6}.$$

Without unbiased assumption, we know that (57) still holds. Similarly, the way to get (46), (62), and (64), we have

$$\begin{aligned} \mathbf{E}_{\check{\delta}_k}[W_{1,k+1}] &\leq W_{1,k} - \|\mathbf{x}_k\|_{\eta\alpha L - \frac{\eta}{2}\mathbf{K} - \frac{3\eta^2\alpha^2}{2}L^2 - \eta(1+3\eta)L_f^2\mathbf{K}}^2 \\ &\quad - \eta\beta\mathbf{x}_k^T\mathbf{K}\left(\mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0\right) + \left\|\mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0\right\|_{\frac{3\eta^2\beta^2}{2}\mathbf{K}}^2 \\ &\quad + \eta(1 + 3\eta)n\sigma^2 \end{aligned} \quad (142a)$$

$$\begin{aligned} \mathbf{E}_{\check{\delta}_k}[W_{3,k+1}] &\leq W_{3,k} - \eta\alpha\mathbf{x}_k^T\mathbf{L}\left(\mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0\right) \\ &\quad + \|\mathbf{x}_k\|_{\eta(\beta L + \frac{1}{2}\mathbf{K}) + \eta^2\left(\frac{\alpha^2}{2} - \alpha\beta + \beta^2\right)L^2 + \eta(1+2\eta)L_f^2\mathbf{K}}^2 \\ &\quad + \frac{\eta}{2\beta^2}(1 + 3\eta)L_f^2\mathbf{E}_{\check{\delta}_k}[\|\check{\mathbf{g}}_k^u\|^2] + \eta(1 + 2\eta)n\sigma^2 \\ &\quad - \left\|\mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0\right\|_{\eta(\beta - \frac{1}{2} - \eta\beta^2)\mathbf{K}}^2 \end{aligned} \quad (142b)$$

$$\begin{aligned} \mathbf{E}_{\check{\delta}_k}[W_{4,k+1}] &\leq W_{4,k} - \frac{\eta}{4}(1 - 2\eta L_f)\mathbf{E}_{\check{\delta}_k}[\|\check{\mathbf{g}}_k^u\|^2] \\ &\quad + \|\mathbf{x}_k\|_{\eta L_f^2\mathbf{K}}^2 - \frac{\eta}{4}\|\check{\mathbf{g}}_k^0\|^2 + n\sigma^2\eta. \end{aligned} \quad (142c)$$

Then, similarly, the way to get (68a), from (57) and (142a)–(142c), we have

$$\begin{aligned} \mathbf{E}_{\check{\delta}_k}[W_{k+1}] &\leq W_k - \|\mathbf{x}_k\|_{\varepsilon_3\mathbf{K}}^2 - \left\|\mathbf{v}_k + \frac{1}{\beta}\mathbf{g}_k^0\right\|_{\varepsilon_4\mathbf{K}}^2 \\ &\quad - \frac{1}{4}\eta\|\check{\mathbf{g}}_k^0\|^2 + \eta(3 + 5\eta)n\sigma^2. \end{aligned} \quad (143)$$

Then, similarly, the way to get (19), from (143) and (90), we have (21). ■

REFERENCES

- [1] J. Dean, G. Corrado, R. Monga, K. Chen, M. Devin, M. Mao, M. Ranzato, A. Senior, P. Tucker, K. Yang, Q. V. Le, and A. Y. Ng, “Large scale distributed deep networks,” in *Advances in Neural Information Processing Systems*, 2012, pp. 1223–1231.
- [2] H. B. McMahan, E. Moore, D. Ramage, *et al.*, “Communication-efficient learning of deep networks from decentralized data,” in *Proc. Int. Conf. Artificial Intelligence and Statistics*, 2017, pp. 1273–1282.
- [3] T. Tatarenko and B. Touri, “Non-convex distributed optimization,” *IEEE Trans. Automatic Control*, vol. 62, no. 8, pp. 3744–3757, 2017.
- [4] M. Hong, D. Hajinezhad, and M.-M. Zhao, “Prox-PDA: The proximal primal-dual algorithm for fast distributed nonconvex optimization and learning over networks,” in *Proc. Int. Conf. Machine Learning*, 2017, pp. 1529–1538.

- [5] A. Daneshmand, G. Scutari, and V. Kungurtsev, "Second-order guarantees of gradient algorithms over networks," in *Proc. Annual Allerton Conf. Communication, Control, and Computing*, 2018, pp. 359–365.
- [6] H. Sun and M. Hong, "Distributed non-convex first-order optimization and information processing: Lower complexity bounds and rate optimal algorithms," arXiv preprint arXiv: 1804.02729, 2018.
- [7] D. Hajinezhad and M. Hong, "Perturbed proximal primal-dual algorithm for nonconvex nonsmooth optimization," *Mathematical Programming*, vol. 176, no. 1, pp. 207–245, 2019.
- [8] H. Sun and M. Hong, "Distributed non-convex first-order optimization and information processing: Lower complexity bounds and rate optimal algorithms," *IEEE Trans. Signal Processing*, vol. 67, no. 22, pp. 5912–5928, 2019.
- [9] H.-T. Wai, J. Lafond, A. Scaglione, and E. Moulines, "Decentralized Frank-Wolfe algorithm for convex and nonconvex problems," *IEEE Trans. Automatic Control*, vol. 62, no. 11, pp. 5522–5537, 2017.
- [10] J. Langford, L. Li, and T. Zhang, "Sparse online learning via truncated gradient," *Journal of Machine Learning Research*, vol. 10, no. 3, pp. 777–801, 2009.
- [11] B. Recht, C. Re, S. Wright, and F. Niu, "Hogwild!: A lock-free approach to parallelizing stochastic gradient descent," in *Advances in Neural Information Processing Systems*, 2011, pp. 693–701.
- [12] C. M. De Sa, C. Zhang, K. Olukotun, and C. Ré, "Taming the wild: A unified analysis of hogwild-style algorithms," in *Advances in Neural Information Processing Systems*, 2015, pp. 2674–2682.
- [13] X. Lian, Y. Huang, Y. Li, and J. Liu, "Asynchronous parallel stochastic gradient for nonconvex optimization," in *Advances in Neural Information Processing Systems*, 2015, pp. 2737–2745.
- [14] X. Lian, H. Zhang, C.-J. Hsieh, Y. Huang, and J. Liu, "A comprehensive linear speedup analysis for asynchronous stochastic parallel optimization from zeroth-order to first-order," in *Advances in Neural Information Processing Systems*, 2016, pp. 3054–3062.
- [15] Z. Zhou, P. Mertikopoulos, N. Bambos, P. Glynn, Y. Ye, L.-J. Li, and F.-F. Li, "Distributed asynchronous optimization with unbounded delays: How slow can you go?" in *Proc. Int. Conf. Machine Learning*, 2018, pp. 5970–5979.
- [16] J. Bernstein, Y.-X. Wang, K. Azizzadenesheli, and A. Anandkumar, "SignSGD: Compressed optimisation for non-convex problems," in *Proc. Int. Conf. Machine Learning*, 2018, pp. 560–569.
- [17] P. Jiang and G. Agrawal, "A linear speedup analysis of distributed deep learning with sparse and quantized communication," in *Advances in Neural Information Processing Systems*, 2018, pp. 2525–2536.
- [18] A. Reiszadeh, A. Mokhtari, H. Hassani, A. Jadbabaie, and R. Pedarsani, "FedPAQ: A communication-efficient federated learning method with periodic averaging and quantization," in *Proc. Int. Conf. Artificial Intelligence and Statistics*, 2020, pp. 2021–2031.
- [19] D. Basu, D. Data, C. Karakus, and S. Diggavi, "Qsparse-local-SGD: Distributed SGD with quantization, sparsification and local computations," in *Advances in Neural Information Processing Systems*, 2019, pp. 14668–14679.
- [20] J. Wang and G. Joshi, "Adaptive communication strategies to achieve the best error-runtime trade-off in local-update SGD," in *Proc. Conf. Machine Learning and Systems*, 2019, pp. 212–229.
- [21] H. Yu, S. Yang, and S. Zhu, "Parallel restarted SGD with faster convergence and less communication: Demystifying why model averaging works for deep learning," in *Proc. AAAI Conf. Artificial Intelligence*, 2019, pp. 5693–5700.
- [22] F. Haddadpour, M. M. Kamani, M. Mahdavi, and V. Cadambe, "Trading redundancy for communication: Speeding up distributed SGD for nonconvex optimization," in *Proc. Int. Conf. Machine Learning*, 2019, pp. 2545–2554.
- [23] H. Yu, R. Jin, and S. Yang, "On the linear speedup analysis of communication efficient momentum SGD for distributed non-convex optimization," in *Proc. Int. Conf. Machine Learning*, 2019, pp. 7184–7193.
- [24] F. Haddadpour, M. M. Kamani, M. Mahdavi, and V. Cadambe, "Local SGD with periodic averaging: Tighter analysis and adaptive synchronization," in *Advances in Neural Information Processing Systems*, 2019, pp. 11080–11092.
- [25] H. Yu and R. Jin, "On the computation and communication complexity of parallel SGD with dynamic batch sizes for stochastic non-convex optimization," in *Proc. Int. Conf. Machine Learning*, 2019, pp. 7174–7183.
- [26] Z. Jiang, A. Balu, C. Hegde, and S. Sarkar, "Collaborative deep learning in fixed topology networks," in *Advances in Neural Information Processing Systems*, 2017, pp. 5904–5914.
- [27] X. Lian, C. Zhang, H. Zhang, C.-J. Hsieh, W. Zhang, and J. Liu, "Can decentralized algorithms outperform centralized algorithms? A case study for decentralized parallel stochastic gradient descent," in *Advances in Neural Information Processing Systems*, 2017, pp. 5330–5340.
- [28] J. George, T. Yang, H. Bai, and P. Gurrarn, "Distributed stochastic gradient method for non-convex problems with applications in supervised learning," in *Proc. IEEE Conf. Decision and Control*, 2019, pp. 5538–5543.
- [29] X. Lian, W. Zhang, C. Zhang, and J. Liu, "Asynchronous decentralized parallel stochastic gradient descent," in *Proc. Int. Conf. Machine Learning*, 2018, pp. 3043–3052.
- [30] M. Assran, N. Loizou, N. Ballas, and M. Rabbat, "Stochastic gradient push for distributed deep learning," in *Proc. Int. Conf. Machine Learning*, 2019, pp. 344–353.
- [31] H. Tang, S. Gan, C. Zhang, T. Zhang, and J. Liu, "Communication compression for decentralized training," in *Advances in Neural Information Processing Systems*, 2018, pp. 7652–7662.
- [32] A. Reiszadeh, H. Taheri, A. Mokhtari, H. Hassani, and R. Pedarsani, "Robust and communication-efficient collaborative learning," in *Advances in Neural Information Processing Systems*, 2019, pp. 8386–8397.
- [33] H. Taheri, A. Mokhtari, H. Hassani, and R. Pedarsani, "Quantized decentralized stochastic learning over directed graphs," in *Proc. Int. Conf. Machine Learning*, 2020, pp. 9324–9333.
- [34] N. Singh, D. Data, J. George, and S. Diggavi, "SQuARM-SGD: Communication-efficient momentum SGD for decentralized optimization," *IEEE Journal on Selected Areas in Information Theory*, vol. 2, no. 3, pp. 954–969, 2021.
- [35] J. Wang and G. Joshi, "Cooperative SGD: A unified framework for the design and analysis of communication-efficient SGD algorithms," *Journal of Machine Learning Research*, vol. 22, no. 213, pp. 1–50, 2021.
- [36] H. Tang, X. Lian, M. Yan, C. Zhang, and J. Liu, "D²: Decentralized training over decentralized data," in *Proc. Int. Conf. Machine Learning*,

- 2018, pp. 4848–4856.
- [37] S. Lu, X. Zhang, H. Sun, and M. Hong, “GNSD: A gradient-tracking based nonconvex stochastic algorithm for decentralized optimization,” in *Proc. IEEE Data Science Workshop*, 2019, pp. 315–321.
- [38] J. Zhang and K. You, “Decentralized stochastic gradient tracking for empirical risk minimization,” arXiv preprint arXiv: 1909.02712, 2019.
- [39] S. U. Stich, “Local SGD converges fast and communicates little,” in *Proc. Int. Conf. Learning Representations*, 2019.
- [40] A. Koloskova, S. Stich, and M. Jaggi, “Decentralized stochastic optimization and gossip algorithms with compressed communication,” in *Proc. Int. Conf. Machine Learning*, 2019, pp. 3478–3487.
- [41] S. Pu, A. Olshevsky, and I. C. Paschalidis, “A sharp estimate on the transient time of distributed stochastic gradient descent,” *IEEE Transactions on Automatic Control*, 2021. DOI: 10.1109/TAC.2021.3126253
- [42] M. Rabbat, “Multi-agent mirror descent for decentralized stochastic optimization,” in *Proc. Int. Workshop on Computational Advances in Multi-Sensor Adaptive Processing*, 2015, pp. 517–520.
- [43] G. Lan, S. Lee, and Y. Zhou, “Communication-efficient algorithms for decentralized and stochastic optimization,” *Mathematical Programming*, vol. 180, no. 1, pp. 237–284, 2020.
- [44] D. Yuan, Y. Hong, D. W. Ho, and G. Jiang, “Optimal distributed stochastic mirror descent for strongly convex optimization,” *Automatica*, vol. 90, pp. 196–203, 2018.
- [45] D. Jakovetic, D. Bajovic, A. K. Sahu, and S. Kar, “Convergence rates for distributed stochastic optimization over random networks,” in *Proc. IEEE Conf. Decision and Control*, 2018, pp. 4238–4245.
- [46] A. Fallah, M. Gurbuzbalaban, A. Ozdaglar, U. Simsekli, and L. Zhu, “Robust distributed accelerated stochastic gradient methods for multiagent networks,” arXiv preprint arXiv: 1910.08701, 2019.
- [47] S. Pu and A. Garcia, “Swarming for faster convergence in stochastic optimization,” *SIAM Journal on Control and Optimization*, vol. 56, no. 4, pp. 2997–3020, 2018.
- [48] S. Pu and A. Nedić, “A distributed stochastic gradient tracking method,” in *Proc. IEEE Conf. Decision and Control*, 2018, pp. 963–968.
- [49] R. Xin, A. K. Sahu, U. A. Khan, and S. Kar, “Distributed stochastic optimization with gradient tracking over strongly-connected networks,” in *Proc. IEEE Conf. Decision and Control*, 2019, pp. 8353–8358.
- [50] M. Mesbahi and M. Egerstedt, *Graph Theoretic Methods in Multiagent Networks*. Princeton University Press, 2010.
- [51] A. Rakhlin, O. Shamir, and K. Sridharan, “Making gradient descent optimal for strongly convex stochastic optimization,” in *Proc. Int. Conf. Machine Learning*, 2012, pp. 1571–1578.
- [52] H. Karimi, J. Nutini, and M. Schmidt, “Linear convergence of gradient and proximal-gradient methods under the Polyak-Lojasiewicz condition,” in *Proc. Joint European Conf. Machine Learning and Knowledge Discovery in Databases*, 2016, pp. 795–811.
- [53] H. Zhang and L. Cheng, “Restricted strong convexity and its applications to convergence analysis of gradient-type methods in convex optimization,” *Optimization Letters*, vol. 9, no. 5, pp. 961–979, 2015.
- [54] Z. Li and J. Li, “A simple proximal stochastic gradient method for nonsmooth nonconvex optimization,” in *Advances in Neural Information Processing Systems*, 2018, pp. 5569–5579.
- [55] M. Fazel, R. Ge, S. Kakade, and M. Mesbahi, “Global convergence of policy gradient methods for the linear quadratic regulator,” in *Proc. Int. Conf. Machine Learning*, 2018, pp. 1467–1476.
- [56] W. Shi, Q. Ling, G. Wu, and W. Yin, “EXTRA: An exact first-order algorithm for decentralized consensus optimization,” *SIAM Journal on Optimization*, vol. 25, no. 2, pp. 944–966, 2015.
- [57] A. Nedić, A. Olshevsky, and W. Shi, “Achieving geometric convergence for distributed optimization over time-varying graphs,” *SIAM Journal on Optimization*, vol. 27, no. 4, pp. 2597–2633, 2017.
- [58] H. Li and Z. Lin, “Revisiting extra for smooth distributed optimization,” *SIAM Journal on Optimization*, vol. 30, no. 3, pp. 1795–1821, 2020.
- [59] Y. LeCun, C. Cortes, and C. Burges, “MNIST handwritten digit database,” [Online] Available: <http://yann.lecun.com/exdb/mnist>, 2010.
- [60] L. Bottou, “Stochastic gradient descent tricks,” in *Neural Networks: Tricks of the Trade*. Springer, 2012, pp. 421–436.
- [61] Y. Nesterov, *Lectures on Convex Optimization*, 2nd ed. Springer Int. Publishing, 2018.
- [62] Y. Tang, J. Zhang, and N. Li, “Distributed zero-order algorithms for nonconvex multiagent optimization,” *IEEE Trans. Control of Network Systems*, vol. 8, no. 1, pp. 269–281, 2020.
- [63] X. Yi, L. Yao, T. Yang, J. George, and K. H. Johansson, “Distributed optimization for second-order multi-agent systems with dynamic eventtriggered communication,” in *Proc. IEEE Conf. Decision and Control*, 2018, pp. 3397–3402.



Xinlei Yi received the Ph.D. degree in electrical engineering from the School of Electrical Engineering and Computer Science, KTH Royal Institute of Technology in 2020 and now is a Postdoc at the same university. He received B.S. and M.S. degrees in mathematics from China University of Geoscience and Fudan University in 2011 and 2014, respectively. His current research interests include online optimization, distributed optimization, and event-triggered control.



Shengjun Zhang received the B.Eng. degree in automation of honors program from China Agricultural University in 2014, and the M.S. degree in electrical engineering from New York University, USA, in 2017. He is currently working toward the Ph.D. degree with the Department of Electrical Engineering in the College of Engineering, University of North Texas, USA. His current research interests include distributed optimization, statistical learning, and Sparse PCA.



Tao Yang is a Professor at the State Key Laboratory of Synthetical Automation for Process Industries, Northeastern University. He was an Assistant Professor at the Department of Electrical Engineering, University of North Texas, USA, from 2016–2019. He received the Ph.D. degree in electrical engineering from Washington State University in 2012. Between August 2012 and August 2014, he was an ACCESS Post-Doctoral Researcher with the ACCESS Linnaeus Centre, Royal Institute of Technology, Sweden. He then joined the Pacific Northwest National Laboratory as a Postdoc, and was promoted to Scientist/Engineer II in 2015. His research interests include industrial artificial intelligence, integrated optimization and control, distributed control and optimization with applications to process industries, cyber-physical systems, and networked control systems. He is an Associate Editor for *IEEE Transactions on Control Systems Technology*, *IEEE Transactions on Neural Networks and Learning*

Systems, and *IEEE/CAA Journal of Automatica Sinica*. He is members of several technical committees of the IEEE Control Systems Society and IFAC. He received Ralph E. Powe Junior Faculty Enhancement Award and Best Student Paper award (as an advisor) of the 14th IEEE International Conference on Control & Automation in 2018.



Tianyou Chai (Fellow, IEEE) received the Ph.D. degree in control theory and engineering in 1985 from Northeastern University where he became a Professor in 1988. He is the Founder and Director of the Center of Automation, which became a National Engineering and Technology Research Center and a State Key Laboratory. He is a Member of Chinese Academy of Engineering, IFAC Fellow, and IEEE Fellow. He has served as Director of Department of Information Science of National Natural Science Foundation of China from 2010 to 2018. His current research interests include modeling, control, optimization and integrated automation of complex industrial processes.

He has published 297 peer reviewed international journal papers. His paper titled Hybrid intelligent control for optimal operation of shaft furnace roasting process was selected as one of three best papers for the Control Engineering Practice Paper Prize for 2011–2013. He has developed control technologies with applications to various industrial processes. For his contributions, he has won 5 prestigious awards of National Natural Science,

National Science and Technology Progress and National Technological Innovation, the 2007 Industry Award for Excellence in Transitional Control Research from IEEE Multiple-conference on Systems and Control, and the 2017 Wook Hyun Kwon Education Award from Asian Control Association.



Karl Henrik Johansson (Fellow, IEEE) is Professor at the School of Electrical Engineering and Computer Science, KTH Royal Institute of Technology. He received M.Sc. and Ph.D. degrees from Lund University. He has held visiting positions at UC Berkeley, Caltech, NTU, HKUST Institute of Advanced Studies, and NTNU. His research interests are in networked control systems, cyber-physical systems, and applications in transportation, energy, and automation. He has served on the IEEE Control Systems Society Board of Governors, the IFAC Executive Board, and the European Control Association Council. He has received several best paper awards and other distinctions from IEEE and ACM. He has been awarded Distinguished Professor with the Swedish Research Council and Wallenberg Scholar with the Knut and Alice Wallenberg Foundation. He has received the Future Research Leader Award from the Swedish Foundation for Strategic Research and the triennial Young Author Prize from IFAC. He is Fellow of the IEEE and the Royal Swedish Academy of Engineering Sciences, and he is IEEE Control Systems Society Distinguished Lecturer.