

An Algebraic Proof of Thurston's Rigidity for Maps With a Superattracting Cycle

Alon Levy

September 23, 2014

Abstract

We study rational self-maps of \mathbb{P}^1 whose critical points all have finite forward orbit. Thurston's rigidity theorem states that outside a single well-understood family, there are finitely many such maps over \mathbb{C} of fixed degree and critical orbit length. We provide an algebraic proof of this fact for tamely ramified maps for which at least one of the critical points is periodic. We also produce wildly ramified counterexamples.

1 Introduction

The behavior of a rational self-map of \mathbb{P}^1 of degree $d \geq 2$ at its critical points plays an important role in its global dynamics. Specifically, we study a special set of maps:

Definition 1.1. A rational map $\varphi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is **postcritically finite** (PCF) if all of its critical points have finite forward orbit.

PCF maps have attracted some attention from complex dynamicists [15, 6], who relate their combinatorial properties to their dynamical properties. At the same time, arithmetic dynamicists have studied their special Galois orbit properties. In brief, if we fix a point $z \in \mathbb{P}^1(K)$ for some arithmetically interesting field K , the absolute Galois group $\text{Gal}(\overline{K}/K)$ will act on the infinite tree of preimages of z . Conjecturally the Galois action has more or less full image, but if φ is PCF, then the image is much smaller [1, 3, 4]. If one views rational maps as analogous to elliptic curves, as is the approach used in [23], then PCF maps are thus somewhat analogous to elliptic curves with complex multiplication.

Our goal in this paper is to study the PCF maps from the point of view of the moduli space of rational maps on \mathbb{P}^1 . We write rational maps by their coordinates,

$$\varphi(z) = \frac{f(z)}{g(z)} = \frac{a_d z^d + \dots + a_0}{b_d z^d + \dots + b_0}$$

The space of morphisms is a subset of the space

$$(a_d : \dots : a_0 : b_d : \dots : b_0) \in \mathbb{P}^{2d+1}$$

defined by the open affine condition that $\gcd(f, g) = 1$; we call this open affine subspace Rat_d . The geometry of φ is preserved under coordinate-change; in particular, if $A \in \text{PGL}(2)$ and φ is PCF, then so is its conjugation $A\varphi A^{-1}$. Thus we need to quotient the space by $\text{PGL}(2)$ -conjugation. We set $\text{Rat}_d // \text{PGL}(2) = \text{M}_d$; as established in [22, 20, 16], the quotient M_d is geometric and the stabilizer groups in $\text{PGL}(2)$ are finite.

In particular, $\dim \text{M}_d = 2d - 2$. Conversely, when φ is tamely ramified it has $2d - 2$ critical points, counted with multiplicity. Although the condition that φ is PCF cannot be expressed as a finite number of equations, if we specify the size of the forward orbit of each critical point, then we obtain an algebraic equation. Hence we obtain $2d - 2$ equations. It is therefore natural to ask if those equations always intersect in a finite number of points.

Unfortunately, in one special case, the $2d - 2$ equations will intersect in a curve, rather than in finitely many points:

Definition 1.2. The map $\varphi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is called a **Lattès map** if there is an elliptic curve E , a morphism $\alpha : E \rightarrow E$, and a finite separable map π such that the following diagram commutes:

$$\begin{array}{ccc} E & \xrightarrow{\alpha} & E \\ \pi \downarrow & & \downarrow \pi \\ \mathbb{P}^1 & \xrightarrow{\varphi} & \mathbb{P}^1 \end{array}$$

If we choose π to be the projection by $P \sim -P$, then α must be of the form $\alpha_{m,T} : P \mapsto mP + T$ where $T \in E[2]$.

The points of a Lattès map $\varphi_{E,\alpha}$ with finite forward image are precisely the points that come from the torsion points on E . Moreover, because α is unramified, the critical points come from the critical points of π , and those are necessarily torsion points; hence, all Lattès maps are PCF. Conversely, Lattès maps $\varphi_{E,\alpha}$ with a fixed (more precisely, continuously varying) α form a curve in M_{m^2} according to the j -invariants of E . For a more complete treatment, see [23].

However, as shown by Thurston, the Lattès maps are the only counterexample to the expectation that the equations defining PCF maps intersect in finitely many points. Thurston studied PCF maps based on underlying combinatorial data:

Definition 1.3. The **critical portrait** of a PCF map φ is a directed graph on its critical points and their forward images in which the edge $x \mapsto y$ occurs if and only if $\varphi(x) = y$. This includes the case in which $x = y$.

All Lattès maps with the same α have the same critical portrait. We have,

Theorem 1.4. (*Thurston's Rigidity* [7, 5]) *Over \mathbb{C} , each critical portrait admits only finitely many non-Lattès maps $\varphi \in \text{M}_d$, which are defined over $\overline{\mathbb{Q}}$.*

It is expected that the scheme defining each portrait is in fact reduced. Epstein shows in Section 5 of [8] that the critical orbit relations, chosen in a specific way to avoid double-counting, intersect transversely (the definition of double-counting is beyond the scope of this paper, but can be thought of as a precise way to say that the critical portraits do not degenerate). Favre and Gauthier [11] show this explicitly for some relations involving PCF polynomials. This can be viewed as an infinitesimal rigidity statement.

The importance of Thurston's rigidity goes beyond PCF maps. Indeed, it is useful when discussing the general behavior of periodic points of φ , that is points z for which $\varphi^n(z) = z$. First, we recall:

Definition 1.5. The **multiplier** of a map φ at a point z of period n , that is a point for which $\varphi^n(z) = z$, is the eigenvalue of the induced action on tangent spaces, $(\varphi^n)'(z)$; it is conjugation-invariant, and invariant for all points in a single periodic cycle.

Definition 1.6. In analogy with the terminology of attracting and repelling periodic points when φ is defined over a valued field, we say a periodic cycle is **superattracting** if its multiplier is 0, an algebraic rather than an analytic condition. Observe that a cycle is superattracting if and only if it contains a critical point.

Epstein uses rigidity to prove a bound on the number of periodic cycles whose multipliers are in the unit disk in [9] and infinitesimal rigidity to locally parametrize rational maps in the last section of [8]. In addition, McMullen [19] uses rigidity to prove a deep structure result on $M_d(\mathbb{C})$:

Theorem 1.7. (*McMullen*) *Let $\Lambda_n : M_d \rightarrow \mathbb{A}^{k_n}$ be the map sending φ to the symmetric functions in the period- n multipliers; here k_n is just the dimension of the target space. For sufficiently large n ,*

$$\Lambda_1 \times \dots \times \Lambda_n : M_d(\mathbb{C}) \rightarrow \mathbb{A}^{k_1 + \dots + k_n}(\mathbb{C})$$

is finite-to-one away from the Lattès curves.

Since the space M_d is defined over \mathbb{Z} , and the PCF locus is defined over \mathbb{Z} as well, it is reasonable to ask if rigidity and McMullen's result can be extended to arithmetically interesting fields. Although they extend to number fields by the Lefschetz principle, the proofs of both rigidity and the derivation of McMullen's result from rigidity employ transcendental techniques, and therefore do not extend to characteristic p . Finding arithmetic analogs of both results is an active research topic in arithmetic dynamics, with some partial results due to Epstein [10], who uses p -adic valuations to prove rigidity for a special class of polynomial maps, and Hutz and Tepper [13], who prove that McMullen's result is true for a generic polynomial over \mathbb{Z} and compute the degree of the finite-to-one map.

Upon looking at various wildly ramified cases, the author's hopes that the two results would be true verbatim in characteristic p were dashed. For example,

if k is a field of characteristic p , then the curve

$$\varphi_t(z) = z^{mp} + tz, t \in k$$

is a non-isotrivial curve in M_{mp} since the multiplier at 0 varies, but is PCF since the only critical point is ∞ and is clearly fixed. In general, if $p < d$ and $p \nmid d$ then $z^d + tz^p$ only has 0 and ∞ for critical points and both are fixed, and can be seen to be a curve in M_d by direct examination of the action of each $A \in \text{PGL}(2)$ on the coefficients.

For a counterexample to a positive-characteristic analog of McMullen, we can take any sufficiently high-dimension family with constant derivative, i.e. $\varphi(z) = cz + f(z^p)/g(z^p)$ where c is a constant, which is again wildly ramified. For this family to not map to a single point in M_d , we require $f(z)/g(z)$ to have degree 2 or higher and thus can be used when $p \mid d$ and $p < d$, but not when $p = d$, and indeed Silverman [22] proved that Λ_1 is an isomorphism from M_2 to \mathbb{A}^2 over \mathbb{Z} .

If we think of rigidity as the statement that $2d - 2$ equations over a space of dimension $2d - 2$ should intersect in finitely many points, then its failure in the wildly ramified case is not surprising, since there are fewer than $2d - 2$ critical points even counted with multiplicity. It is therefore reasonable to propose,

Conjecture 1.8. Over $\overline{\mathbb{F}_p}$, each critical portrait admits only finitely many non-Lattès tamely ramified maps $\varphi \in M_d$.

In positive characteristic, transversality fails, even in large characteristic. Indeed, for quadratic polynomials of the form $z^2 + c$, the critical point is 0, and the condition that the critical point has exact period 3 is $c^3 + 2c^2 + c + 1$, a polynomial of discriminant -23 . It is an open question whether, just for quadratic polynomials, the set of primes for which transversality fails has zero density. For our purposes the question then is whether we can prove finiteness, and in the sequel, we will take rigidity to mean the finiteness portion of Thurston's rigidity.

It is currently beyond the author's means to prove Conjecture 1.8. The proof of rigidity in [7] is topological, and has portions that cannot yet be generalized beyond \mathbb{C} . However, in some cases, a direct algebraic proof is feasible. The case of polynomials is already known:

Theorem 1.9. [2] *Conjecture 1.8 holds for polynomials whenever whenever their reduction after any coordinate change is still tamely ramified, which is the case when they are compositions of polynomials of degree less than p . If the reduction is bad, the reduction must be tamely ramified after clearing common roots of the numerator and denominator.*

By reduction and coordinate change, we mean the following:

Definition 1.10. Let K be a complete algebraically closed non-archimedean valued field, with ring of integers \mathcal{O}_K and maximal ideal \mathfrak{m}_K . Let $\varphi : \mathbb{P}_K^1 \rightarrow \mathbb{P}_K^1$. We call $\psi(z) : \mathbb{P}_K^1 \rightarrow \mathbb{P}_K^1$ an **integral model** for φ if we take the coordinates

of ψ to lie in \mathcal{O}_K and if there exists $A \in \mathrm{PGL}(2, K)$ such that $\varphi = A\psi A^{-1}$, or, equivalently, if ψ and φ reduce to the same point in $\mathrm{M}_d(K)$. We say that φ has **good reduction** if there exists an integral model

$$\psi(z) = \frac{f(z)}{g(z)} = \frac{a_d z^d + \dots + a_0}{b_d z^d + \dots + b_0}$$

such that when we reduce all coefficients mod \mathfrak{m}_K , we have, denoting reduction with overlines,

$$\overline{\psi}(z) = \frac{\overline{f}(z)}{\overline{g}(z)} = \frac{\overline{a_d} z^d + \dots + \overline{a_0}}{\overline{b_d} z^d + \dots + \overline{b_0}} \in \mathrm{Rat}_d(\mathcal{O}_K/\mathfrak{m}_K)$$

i.e. $\overline{f}(z)$ and $\overline{g}(z)$ do not have common roots. Equivalently, φ has good reduction if it reduces to an \mathcal{O}_K -point of the affine space $\mathrm{M}_d(K)$.

Remark 1.11. By a theorem of Hsia [12], φ has good reduction (over an algebraically closed field) only if every multiplier of every periodic cycle lies in \mathcal{O}_K ; the converse is not true.

Ingram argues in private communication that a modification of the argument in [14] could produce a second proof of Theorem 1.9 over a finite field of characteristic $p > d$ using his theory of critical heights of polynomials.

We will prove a stronger result, using purely algebraic methods:

Theorem 1.12. *Conjecture 1.8 for tamely ramified rational maps with a superattracting cycle.*

In Section 2, we will prove Theorem 1.12 by looking at a grand coordinate ring that encodes not only the coefficients a_i and b_i of each rational map φ satisfying the hypotheses of the theorem, but also the critical points and the critical values. We turn this into a graded ring by assigning each critical point weight 1 and assigning other variables weights that make the defining equations for φ homogeneous. We then show that instead of looking at the system of equations $\varphi^n(\zeta_i) = \varphi^m(\zeta_i)$ for all critical points ζ_i (here, n and m are fixed integers, and for fixed n and m we have just finitely many critical portraits), we can look only at the top-degree portions of those equations, which are much simpler. A series of reductions then converts rigidity into a problem of parametrizing rational maps by their critical values, a solved problem using the theory of branched covers.

What we actually prove is not a statement about geometric loci of PCF maps, but about the corresponding rings. We will actually prove that if we take a carefully chosen slice X of Rat_d with ∞ a fixed critical point such that X maps finite-to-one into M_d , then a coordinate ring that is a finite extension of \mathcal{O}_X modulo the PCF ideal $\varphi^n(\zeta_i) = \varphi^m(\zeta_i)$ yields a ring with only finitely many possibilities for the critical value set. Thus there are only finitely many tamely ramified morphisms in the ring.

This method naturally fails in the two known exceptions for rigidity: it fails in the Lattès case because the Lattès maps' critical points are preperiodic but

never periodic, and it fails in wildly ramified cases because multiple parts of the proof require tame ramification for finiteness to work. A full generalization to tamely ramified non-Lattès maps is unlikely. Unless we can produce a critical point that maps to itself under some iterate of φ , the equations $\varphi^n(\zeta_i) = \varphi^m(\zeta_i)$ are already homogeneous with respect to the grading we use, and there is no hope of simplifying them.

We also investigate integrality, a related result, first investigated by Epstein in [10] as a way of proving rigidity for polynomials of prime-power degree. Instead of proving the finiteness of a module over a field k , we can prove the finiteness of a module over a ring, say a local ring \mathcal{O}_K . This turns out to be equivalent to the statement that the set of critical values of a PCF map is integral. In Section 3, we prove,

Corollary 1.13. *Let $\varphi(z)$ be a PCF map with a superattracting cycle defined over a local field, such that any integral model of $\varphi(z)$ has tamely ramified reduction. Then the configuration of the critical values of $\varphi(z)$ is integral.*

Only this definition of integrality is true. A PCF map satisfying the hypotheses of Corollary 1.13 does not have to be integral itself, i.e. have good reduction; only its critical value set must be integral. In personal communication Rivera-Letelier provided a PCF map satisfying the hypothesis of Corollary 1.13 that nonetheless has bad reduction (Example 3.2).

A separate approach to proving rigidity is to use McMullen's theorem. McMullen's proof is also transcendental, but it appears slightly more amenable to attacking in positive characteristic using non-archimedean analysis. We also have,

Theorem 1.14. *[2] Let K be a function field of characteristic p . If φ is PCF and is the composition of maps of degree less than p then its multipliers lie in the field of constants; in other words, McMullen implies rigidity.*

Since in [22] it is shown that Λ_1 is an isomorphism over \mathbb{Z} between M_2 and a plane in \mathbb{A}^3 , we immediately obtain,

Corollary 1.15. *Thurston's rigidity is true in M_2 in any characteristic except 2.*

Theorem 1.14 is true slightly more generally: if $p \leq d$, it holds as long as φ satisfies the same condition on tamely ramified reductions as in Corollary 1.13. Unfortunately, even this approach is problematic as a means of proving rigidity, since it seems easier to prove rigidity directly than to prove it via McMullen. The results of Theorem 1.12 and Corollary 1.15 remain the strongest positive-characteristic rigidity results known to the author. The work of Pink [21] provides another avenue of proof, but so far is only strong enough to prove Corollary 1.15 as well.

Remark 1.16. Contrary to initial expectations, Theorem 1.12 is not helpful in proving an arithmetic analog of Theorem 1.7. In [17], we prove a partial generalization of Theorem 1.7 to large positive characteristic using nonarchimedean-

analytic methods; Theorem 1.14 provides a key ingredient, but there does not appear to be a way to provide a full generalization even assuming Conjecture 1.8.

2 Finiteness

First, we reduce Theorem 1.12 to a result about maps with a superattracting point:

Proposition 2.1. *Suppose that $C \subset M_d$ is a nontrivial family of rational maps and that $n \geq 1$. Under the iteration morphism \circ^n from M_d to M_{d^n} , $\varphi \mapsto \varphi^n$, the family C is mapped to a nontrivial family.*

Proof. This appears to be well-known to complex and arithmetic dynamicists, but not proven in the literature. We give an elementary proof. Suppose that \circ^n maps C to a single point of M_{d^n} . The configuration of fixed points of $\circ^n(C)$ in \mathbb{P}^1 is clearly a single point in the space of points on \mathbb{P}^1 . Since fixed points of $\varphi \in C$ are also fixed points of φ^n , this implies the configuration of fixed points of $\varphi \in C$ is itself constant (in fact, so is the configuration of period-dividing- n points). Let us now lift C from M_d to Rat_d . If maps in C have at least two distinct fixed points, then we can choose a slice of the pullback of C to Rat_d in which 0 and ∞ are fixed. We write

$$\varphi(z) = \frac{f(z)}{g(z)}, f(z) = \sum_{i=0}^d a_i z^i, g(z) = \sum_{i=0}^d b_i z^i$$

In our parametrization, $a_0 = b_d = 0$. We further fix a slice of the pullback such that $a_d = b_0$. Now the slice has dimension 1 and surjects onto C , and its points are completely determined by roots of $f(z)$ and $g(z)$. But if $\circ^n(C)$ is a point then the configuration of points mapping to each of the fixed points 0 and ∞ is again constant, and if $\varphi(\alpha) = 0$ then $\varphi^n(\alpha) = 0$ and a similar result holds for ∞ .

Finally, suppose maps in C have a unique fixed point, which, in the pullback to Rat_d , we take to be ∞ . Then they may be written as

$$\varphi(z) = z + \frac{1}{g(z)} = \frac{1 + zg(z)}{g(z)}, \deg g = d - 1$$

Furthermore, we can take $g(z)$ to be monic. Again, if $\varphi(\alpha) = \infty$ then $\varphi^n(\alpha) = \infty$, so if $\circ^n(C)$ is a point then we fix the roots of g for all maps $\varphi \in C$, and this again forces C to be a point. \square

Corollary 2.2. *It suffices to prove Theorem 1.12 for maps with a superattracting fixed point.*

Proof. The composition of tamely ramified morphisms is tamely ramified, since the local degree of $\varphi \circ \psi$ at z equals the local degree of ψ at z times the local degree of φ at $\psi(z)$. Now, if C is a nontrivial tamely ramified PCF family with

a superattracting cycle of period n , then by Proposition 2.1, after iteration we obtain a nontrivial tamely ramified PCF family with a superattracting fixed point. \square

In the sequel, we write φ for a rational function of degree d with a multiplicity- e pole at ∞ , where $e > 1$. As in the proof of Proposition 2.1, we write φ as f/g , with $f = \sum a_i z^i$ and $g = \sum b_i z^i$. We assume 0 is a fixed point, i.e. $a_0 = 0$. We lose nothing by making this assumption, since φ has more than one fixed point whenever $e > 1$ (see for example the parametrization in the proof of Proposition 2.1 of maps with a unique fixed point; to ensure $\deg \varphi = d$ we must have $\deg g = d - 1$ and then $e = 1$, a contradiction). We may make one final conjugation, which we use to assume $a_d = b_{d-e}$; we normalize to assume $a_d = b_{d-e} = 1$, which we may do since the diagonal group $T \subset \mathrm{PGL}(2)$ acts on a_d with weight $1 - d$ and on b_{d-e} with weight $-(d - e) \neq 1 - d$. We also assume that the characteristic does not divide any of the ramification degrees. With the above assumptions that $a_0 = 0, a_d = b_{d-e} = 1$, we call the resulting space of maps with a degree- e pole at infinity $X \subseteq \mathrm{Rat}_d$. Note that X maps finite-to-one into M_d .

Lemma 2.3. *In X , the configuration of critical values of φ is enough to specify each tamely ramified φ up to finitely many choices.*

Proof. This follows from standard results on (arbitrary-characteristic) monodromy theory. See [24] for more details. In brief, the tame absolute Galois group of \mathbb{P}^1 minus n points, $\pi_1^{p'}$, is the profinite prime-to- p completion of the free group on $n - 1$ generators (Theorem 4.9.1), and finite unramified covers correspond to finite sets with a continuous left-action of the absolute Galois group (Theorem 4.6.4); for tame ramification, we also require the inertia group at each generator to be prime to p . Now specifying the list of critical values with multiplicities bounds the degree of the cover, bounding the size of those finite sets, and there are finitely many continuous actions of a topologically finitely generated group on a finite set.

Covers of \mathbb{P}^1 are only unique up to the right-action of $\mathrm{PGL}(2)$. While this is not the same as the conjugation action we use to obtain M_d , X maps finite-to-one into the quotient by the right-action as well. This is because we may pick ∞ to be an order- e pole and 0 to be a zero, involving just finitely many choices, and then up to a finite quotient the only $\mathrm{PGL}(2)$ right-action that preserves this choice is the diagonal action. But now the diagonal action of multiplication by t acts on a_d/b_{d-e} as multiplication by t^{-e} , and so we again get just finitely many choices that preserve X . \square

On the affine variety X we would like to show that the equations defining a PCF locus intersect at finitely many points. This is equivalent to showing that the associated coordinate ring is finite as a k -module. More precisely, let ζ_i for $i = 1, \dots, 2d - 1 - e$ be the critical points and ξ_i be the associated critical values. We have simple relations between the critical points and the coordinates a_i, b_i coming from the polynomial in terms of a_i and b_i whose roots

are the critical points, as well as simple relations between the critical points, the coordinates, and the critical values coming from the equation $\varphi(\zeta_i) = \xi_i$. However, we cannot easily pass from coordinates to critical values without going through critical points, requiring us to relate all three quantities.

We use graded rings to relate critical points, critical values, and coordinates. We recall the basic properties of graded rings from Section 13 of [18]. If R is a ring, then a decomposition of the form

$$R = \bigoplus_{i=0}^{\infty} R_i, \quad R_i R_j \subseteq R_{i+j}$$

is called a **grading**, and a ring with a grading is called a **graded ring**. Elements of the summands R_i are called **homogeneous**, and i is called their **degree**; if $x \in R$ is not homogeneous then we say the degree of x is D if $x = x_0 + \dots + x_D$ where $x_i \in R_i$ and $x_D \neq 0$. An ideal $I \triangleleft R$ is called homogeneous if I is generated by homogeneous elements, or, equivalently, if whenever a_1, \dots, a_n are homogeneous and $a_1 + \dots + a_n \in I$, we also have $a_1, \dots, a_n \in I$. If I is homogeneous, then R/I is naturally a graded ring with

$$R/I = \bigoplus_{i=0}^{\infty} R_i / (I \cap R_i)$$

where the quotient on the right-hand side is a quotient of abelian groups.

Now let

$$R = k[\zeta_1, \dots, \zeta_{2d-1-e}, \xi_1, \dots, \xi_{2d-1-e}, a_{d-1}, \dots, a_1, b_{d-e-1}, \dots, b_0]$$

We grade it by setting

$$\deg \zeta_i = 1, \deg \xi_i = e, \deg a_i = d - i, \deg b_i = d - e - i$$

This requires us to mark the critical points, but we lose nothing from doing so, since there are only finitely many of them, which does not change any finiteness question.

We impose the usual relations of coordinates, critical points, and critical values; we call the ideal they generate I_Y and the quotient ring \mathcal{O}_Y . The variety $Y = \text{Spec } \mathcal{O}_Y$ is an S_{2d-1-e} -cover of a variety that contains X as an open dense subvariety but also has degenerate maps, i.e. maps $\varphi = f/g$ where f and g have a common root.

Lemma 2.4. *The ring \mathcal{O}_Y is also graded with the grading naturally obtained from R .*

Proof. It suffices to show that I_Y is generated by homogeneous polynomials relative to the grading that we chose. Now the relations between coordinates and critical points are encoded by the critical point polynomial

$$\sum_{(i,j)=(1,0)}^{(d,d-e)} (i-j)a_i b_j z^{i+j-1} = e \prod_{i=1}^{2d-1-e} (z - \zeta_i)$$

which imposes $2d - 1 - e$ equations, one for each coefficient except the leading term. Now on the left, the z^r -coefficient is the sum of $a_i b_j$'s for which $i + j = r + 1$ and thus its degree is $2d - e - r - 1$, while on the right it is

$$e\sigma_{2d-e-r-1} = e \sum_{l_1 < \dots < l_{2d-e-r-1}} \zeta_{l_1} \cdots \zeta_{l_{2d-e-r-1}}$$

which again has degree $2d - e - r - 1$. Since we assume tame ramification throughout, $e \neq 0$ as an element of the field k , and the right-hand side is not zero. We're now equating homogeneous algebraic expressions of the same degree, so we obtain homogeneous equations.

The relations between critical points and critical values are of the form $\varphi(\zeta_i) = \xi_i$. By symmetry, it suffices to show that the relation is homogeneous for ζ_1 and ξ_1 . Clearing denominators, we obtain $f(\zeta_1) = \xi_1 g(\zeta_1)$. Since $\deg \xi_1 = e$, it suffices to show $f(\zeta_1)$ is homogeneous of degree d and $g(\zeta_1)$ is homogeneous of degree $d - e$. But now each $a_i \zeta_1^i$ term has degree d and each $b_j \zeta_1^j$ term has degree $d - e$, and we are done. \square

Lemma 2.3 simply says that if

$$S = k[\xi_1, \dots, \xi_{2d-1-e}]$$

with a natural inclusion $S \rightarrow R$, then there is a finite module basis for \mathcal{O}_Y that generates all well-defined tamely ramified morphisms in \mathcal{O}_Y over $S/I_Y \cap S$. This is false if we drop either of the conditions: wildly ramified maps have fewer critical points so there are positive-dimension families with constant critical values, for example $z^p + tz$ is a non-isotrivial family whose only critical point and value is ∞ ; in addition, if we allow the maps to degenerate, then each common root of f and g is a critical point and any point in \mathbb{P}^1 can be set as its corresponding critical value without violating the equations defining Y .

We will also use,

Definition 2.5. Let $h \in R$ or $h \in \mathcal{O}_Y$. If $\deg h = D$, we say the **top-degree term** of h is its degree- D part. In particular, it is not necessarily a monomial. In the decomposition

$$R = \bigoplus_{i=0}^{\infty} R_i, \quad \deg x = i \text{ if } x \in R_i$$

we may write $h = h_0 + \dots + h_D$ with $h_i \in R_i$ and then the top-degree term is h_D ; the construction in \mathcal{O}_Y is similar.

The equations defining a PCF locus, $\varphi^m(\zeta_i) = \varphi^n(\zeta_i)$ for all i , become much simpler upon passing to top-degree terms. While each ξ_i is of pure degree e by definition, and by Lemma 2.4 this definition is compatible with the usual equations relating critical points and critical values, $\varphi^n(\zeta_i)$ with $n > 1$ is no longer homogeneous. But as we will see, the assumption that ∞ is a fixed critical

point ensures that the expression $\varphi^n(\zeta_i)$ has degree e^n with a particularly simple top-degree term.

More precisely, write φ^n as f_n/g_n , such that

$$f_n(z) = \sum a_i(f_{n-1}(z))^i(g_{n-1}(z))^{d-i}$$

$$\text{and } g_n(z) = \sum b_i(f_{n-1}(z))^i(g_{n-1}(z))^{d-i}$$

By convention, $f_1 = f$ and $g_1 = g$, or alternatively $f_0(z) = z$ and $g_0(z) = 1$. Then we have:

Lemma 2.6. *For each expression $h \in R$ such that $\deg h = u > 1$, we have $\deg f_n(h) = d^n u$ and $\deg g_n(h) = (d^n - e^n)u$. If the top-degree term of h is equal to h_u , then the top-degree term of $f_n(h)$ is $h_u^{d^n}$ and the top-degree term of $g_n(h)$ is $h_u^{d^n - e^n}$.*

Proof. We use induction on n . If $n = 1$, then $f_n = f$ and $g_n = g$. In the sum

$$\sum_{i=1}^d a_i h^i$$

the degree of the i th term is $d - i + iu$ and since $u > 1$, this degree is strictly maximized when $i = d$ and the degree is equal to du . Likewise, in the sum

$$\sum_{i=0}^{d-e} b_i h^i$$

the degree of the i th term is $d - e - i + iu$ and this is maximized when $i = d - e$ and the degree is equal to $(d - e)u$. In both cases, letting h_u be the top-degree term of h , the top-degree term of $f(h)$ is the top-degree term of h^d , which is just h_u^d , and likewise the top-degree term of $g(h)$ is h_u^{d-e} .

Now, suppose the lemma is true up to $n - 1$. Then in the sum

$$\sum_{i=1}^d a_i (f_{n-1}(h))^i (g_{n-1}(h))^{d-i}$$

the i th term has degree

$$d - i + i \deg f_{n-1}(h) + (d - i) \deg g_{n-1}(h)$$

$$= d - i + i d^{n-1} u + (d - i)(d^{n-1} - e^{n-1})u$$

$$= d + d(d^{n-1} - e^{n-1})u + i(e^{n-1}u - 1)$$

This is (strictly) maximized when i is maximized, i.e. when $i = d$ and the degree is $d^n u$. Thus the top-degree term comes only from $(f_{n-1}(h))^d$, and by the induction hypothesis it is equal to $h_u^{(d^{n-1})d}$.

In the denominator, in the sum

$$\sum_{i=0}^{d-e} b_i (f_{n-1}(h))^i (g_{n-1}(h))^{d-i}$$

the degree of the i th term is $d - e - i + i \deg f_{n-1}(h) + (d - i) \deg g_{n-1}(h)$ which is also maximized when i is maximized, i.e. when $i = d - e$, and then the degree is $(d^n - e^n)u$. As the top-degree term comes only from $(f_{n-1}(h))^{d-e} (g_{n-1}(h))^e$, by the induction hypothesis it is equal to $h_u^{d^{n-1}(d-e) + (d^{n-1} - e^{n-1})e}$, which simplifies to $h_u^{d^n - e^n}$ as required. \square

Now, we will compute the degree of the defining equations $\varphi^m(\zeta_i) = \varphi^n(\zeta_i)$:

Lemma 2.7. *Suppose without loss of generality that $n > m$. Then the equation $\varphi^m(\zeta_i) = \varphi^n(\zeta_i)$ has degree $d^{n-1} + d^{m-1} - e^{m-1}$. Moreover, the top-degree term is $\xi_i^{d^{n-1} + d^{m-1} - e^{m-1}}$.*

Proof. First, since $\varphi(\zeta_i) = \xi_i$, we rewrite the equation as $\varphi^{m-1}(\xi_i) = \varphi^{n-1}(\xi_i)$. Now, we clear denominators and write the equation as

$$f_{m-1}(\xi_i)g_{n-1}(\xi_i) = f_{n-1}(\xi_i)g_{m-1}(\xi_i)$$

Since $\deg \xi_i = e$, the left-hand side has degree $e(d^{m-1} + d^{n-1} - e^{n-1})$ and the right-hand side has degree $e(d^{m-1} + d^{n-1} - e^{m-1})$. Since we assume $n > m$, the top-degree term comes entirely from the right-hand side, and is equal to $\xi_i^{d^{n-1} + d^{m-1} - e^{m-1}}$ since ξ_i is already homogeneous. \square

We will use Lemma 2.7 to show that the defining equations $\varphi^m(\zeta_i) = \varphi^n(\zeta_i)$ intersect in finitely many points (in X) by showing this on the level of top-degree terms. This is a higher-dimensional analog of the fact that a polynomial in one variable with unknown coefficients has finitely many solutions if and only if its top-degree term is nonzero. Theorem 1.12 claims finiteness on the level of a geometric locus, which is equivalent to showing finiteness on the level of rings and modules:

Theorem 2.8. *Let $M = \mathcal{O}_Y/I_{m,n}$, where the ideal $I_{m,n}$ is generated by $\varphi^m(\zeta_i) = \varphi^n(\zeta_i)$ for all i . Then the image ring M' of the map from S to M via R and \mathcal{O}_Y is a finite k -module.*

Proof. It suffices to show that there exists a finite k -submodule of R , which by abuse of notation we also call M , such that each $h \in R$ can be written as $s + r$ where $s \in I_{m,n}$ and $r \in M$. Since the k -submodule

$$\{x \in R : \deg x \leq D\}$$

is finite for each integer D , this is equivalent to proving that for some D , each $h \in R$ can be written as $s + r$ where $s \in I_{m,n}$ and $\deg r \leq D$. We will set $D = d^{n-1} + d^{m-1} - e^{m-1}$, the degree obtained in Lemma 2.7.

Showing that h can be written as $s + r$ with $\deg r \leq D$ is equivalent to showing that whenever $\deg h > D$, we can write h as $s + h_1$ with $\deg h_1 < \deg h$. In one direction the equivalence is obvious; in the other direction, if $h = s + h_1$ and $\deg h_1 > D$ then we can write h_1 as $s_1 + h_2$ where $s_1 \in I_{m,n}$ and $\deg h_2 < \deg h_1$, and after finitely many steps we will obtain h_i such that $\deg h_i \leq D$.

The equation $h = s + h_1$ can also be viewed in terms of top-degree terms. By definition, $\deg h_1 < \deg h$, and so h_1 has no contribution to the top-degree term of the equation. Any lower-degree term in h or s can also be added back to h_1 , and therefore if the equation is satisfied for top-degree terms only, it can be satisfied in general. Put another way, it suffices to show that given a homogeneous h of degree more than D , we can find $s \in I_{m,n}$ such that the top-degree term of s is exactly h (the defining equations for $I_{m,n}$ are not homogeneous, as we will see below).

We replace $I_{m,n}$ with $I'_{m,n}$, whose generators are the top-degree terms of the generators of $I_{m,n}$. It suffices to show that h lies in this ideal: to see why, write the generators of $I_{m,n}$ as s_1, \dots, s_{2d-1-e} , and write their top-degree terms as s'_1, \dots, s'_{2d-1-e} , so that $I'_{m,n} = (s'_1, \dots, s'_{2d-1-e})$. If $h \in I'_{m,n}$, then we can find homogeneous elements in \mathcal{O}_Y , t_1, \dots, t_{2d-1-e} , such that $h = \sum s'_i t_i$, and then $\sum s_i t_i$ has top-degree term equal to h .

Observe now that we can work backward: showing that $h \in I'_{m,n}$ whenever h is homogeneous of degree more than D is equivalent to showing that $\mathcal{O}_Y/I'_{m,n}$ is a finite module. This is because, since $I'_{m,n}$ and h are both homogeneous, the $h = s + r$ condition reduces to $h = s$. Likewise, we can work backward and use the fact that the map $Y \rightarrow X$ is finite to show only that there are finitely many tamely ramified morphisms $\varphi \in Y$ satisfying the equations of $I'_{m,n}$.

By Lemma 2.7, the top-degree term of $\varphi^m(\zeta_i) = \varphi^n(\zeta_i)$ is ξ_i^D . In other words, the generators of $I'_{m,n}$ are ξ_i^D over all i . Now

$$M' = S/(S \cap I_Y, \xi_1^D, \dots, \xi_{2d-e-1}^D)$$

and since

$$S = k[\xi_1, \dots, \xi_{2d-e-1}]$$

this is manifestly a finite k -module. \square

3 Integrality and Critical Integrality

The methods used to prove Theorem 2.8 can also be used to prove an integrality result:

Corollary 3.1. *In Section 2, we can work over \mathbb{Z} and obtain a finite \mathbb{Z} -module in the statement of Theorem 2.8. In other words, if we redefine R as*

$$\mathbb{Z}[\zeta_1, \dots, \zeta_{2d-1-e}, \xi_1, \dots, \xi_{2d-1-e}, a_{d-1}, \dots, a_1, b_{d-e-1}, \dots, b_0]$$

and S as

$$\mathbb{Z}[\xi_1, \dots, \xi_{2d-1-e}]$$

without changing the grading or the defining generators for I_Y and $I_{m,n}$, then the module M' is a finite \mathbb{Z} -module.

Proof. Nowhere in the proof of Theorem 2.8 do we invert elements of k . Therefore, we get

$$M' = \mathbb{Z}[\xi_1, \dots, \xi_{2d-e-1}] / (\xi_1^D, \dots, \xi_{2d-e-1}^D)$$

which is finite as a \mathbb{Z} -module. \square

On less than careful reading, it may appear as if Corollary 3.1 implies that all PCF maps with a superattracting cycle are p -integral, i.e. have good reduction mod p , wherever their reduction (good or bad!) is p -tamely ramified (clearing common roots of \bar{f} and \bar{g} if the reduction is bad). However, in personal communication Rivera-Letelier gave,

Example 3.2. The map

$$\varphi(z) = -45 \frac{3z + 5}{z^2(z - 9)}$$

is PCF but has two 5-adically repelling fixed points, of 5-adic absolute values 1 and $1/5$. Since this map's degree is only 3, this cannot possibly come from wild ramification in the reduction.

In fact, we can instead conclude a weaker result, regarding the integrality of the critical values. But first,

Definition 3.3. Let φ be defined over a local field K . We say that φ is **tame** if, for every integral model of φ over \mathcal{O}_K , the reduction mod the maximal ideal is tamely ramified after clearing common factors of \bar{f} and \bar{g} . If φ is defined over a global field, we say it is tame at a place if it is tame when we regard it as a map over the completion; we say it is tame in general, without specifying a place, if it is tame at every non-archimedean place.

Remark 3.4. An equivalent definition for tameness is that the local degree at each \mathbb{P}^1 -disk is not divisible by the residue characteristic of K . In particular, since composing maps multiplies local degrees, the composition of two tame maps, in particular the iterate of any tame map, is again tame.

When φ is tame, we can talk about configurations of points more freely, since under any $\mathrm{PGL}(2, K)$ -conjugate the map still has tamely ramified reduction. Of course this reduction may be as bad as a degenerate constant map, but it will still have $2d - 2$ critical points and values that we can map the critical points and values of φ to. Using Definition 3.3, we can obtain a more precise formulation of Corollary 1.13:

Corollary 3.5. *Let $\varphi(z)$ be PCF, defined over a local field, tame, and with a superattracting cycle. Then the reduction of φ modulo the maximal ideal has an integral configuration of critical values, i.e. there exists an integral model of φ whose critical values are integral.*

Proof. Let K be the local field of definition and \mathcal{O}_K its ring of integers. Then $\varphi \in M' \otimes \mathcal{O}_K$, a finite \mathcal{O}_K -module. Thus the critical values of φ , i.e. $\xi_1, \dots, \xi_{2d-e-1}$, define a finite \mathcal{O}_K -module, and in particular are \mathcal{O}_K -integral. \square

Remark 3.6. As with Definition 3.3, Corollary 3.5 also applies to maps defined over global fields.

In Lemma 2.3, a configuration of critical values in characteristic p may not correspond to a cover of \mathbb{P}^1 because the lift of a tamely ramified cover of \mathbb{P}^1 to characteristic 0 may have bad reduction; on the level of groups, the tame fundamental group is the prime-to- p profinite completion of the free group on $\#\{\xi_1, \dots, \xi_{2d-e-1}\}$ generators, whereas the characteristic-0 fundamental group is the full profinite completion, which has more quotients and thus more actions. The interpretation of the interplay between Corollary 3.5 and the existence of non-integral PCF maps as in Example 3.2 is that those maps are precisely the bad reduction lifts that prevent Riemann existence from holding verbatim in characteristic p .

For the same reason, we cannot count PCF maps in characteristic p in the same way as maps in characteristic 0. In our case of interest—that is, tamely ramified maps with a superattracting cycle—we can lift each characteristic p PCF configuration of critical values to characteristic 0 and then lift all characteristic p PCF maps, but in the opposite direction we may encounter bad reduction. In particular, there are fewer characteristic p PCF maps than characteristic 0 ones.

The situation for polynomials is much simpler. Not only do we not need S in the proof of Theorem 2.8, but also we obtain:

Theorem 3.7. *Let $f(z)$ be a tame PCF polynomial. Then f has good reduction.*

Proof. See Theorem 7.1 in [2]. The theorem’s statement is weaker, assuming that f is the composition of maps of degree $d < p$, but this assumption is only used to establish that the local degrees on certain disks are never divisible by p , and for that it suffices to assume absolute tame ramification. \square

References

- [1] Wayne Aitken, Farshid Hajir, and Christian Maire, *Finitely ramified iterated extensions*, Int. Math. Res. Not. **14** (2005), 855–880.
- [2] Robert Benedetto, Patrick Ingram, Rafe Jones, and Alon Levy, *Attracting cycles in p -adic dynamics and height bounds for post-critically finite maps*, arXiv:1201.1605v2, to appear in Duke Math. J., Oct 2013.
- [3] Nigel Boston and Rafe Jones, *Arboreal Galois representations*, Geom. Dedicata **124** (2007), no. 1, 27–35.
- [4] ———, *The image of an arboreal Galois representation*, Pure and Applied Mathematics Quarterly **5** (2009), no. 1, 213–225.

- [5] Eva Brezin, Rosemary Byrne, Joshua Levy, Kevin Pilgrim, and Kelly Plummer, *A census of rational maps*, Conformal Geometry and Dynamics **4** (2000), 35–74.
- [6] Xavier Buff, Adam Epstein, Sarah Koch, and Kevin Pilgrim, *On Thurston’s pullback map*, Complex Dynamics: Families and Friends (Dierk Schleicher, ed.), A K Peters, Wellesley, MA, 2009, pp. 561–583.
- [7] Adrien Douady and John H. Hubbard, *A proof of Thurston’s topological characterization of rational functions*, Acta Math. **171** (1993), 263–297.
- [8] Adam Epstein, *Transversality in holomorphic dynamics*, <http://homepages.warwick.ac.uk/~mases/Transversality.pdf>.
- [9] ———, *Infinitesimal Thurston rigidity and the Fatou-Shishikura inequality*, arXiv:math/9902158v1, 1999.
- [10] ———, *Integrality and rigidity for postcritically finite polynomials*, arXiv:1010.2780, 2010.
- [11] Charles Favre and Thomas Gauthier, *Distribution of postcritically finite polynomials*, arXiv:1302.0810, 2013.
- [12] Liang-Chung Hsia, *A weak Néron model with applications to p -adic dynamical systems*, Compositio Math. **100** (1996), 277–304.
- [13] Benjamin Hutz and Michael Tepper, *Multiplier spectra and the moduli space of degree 3 morphisms on \mathbb{P}^1* , arXiv:1110.5082, Oct 2011.
- [14] Patrick Ingram, *A finiteness result for post-critically finite polynomials*, arXiv:1010.3393v2; to appear in Int. Math. Res. Not., Feb 2011.
- [15] Sarah Koch, *Teichmüller theory and critically finite endomorphisms*, Adv. Math. **248** (2013), 573–617.
- [16] Alon Levy, *The space of morphisms on projective space*, Acta Arith. **146** (2011), 13–31.
- [17] ———, *The McMullen map in positive characteristic*, arXiv:1304.2834, Apr 2013.
- [18] Hideyuki Matsumura, *Commutative ring theory*, Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, Cambridge, UK, 1989.
- [19] Curtis T. McMullen, *Families of rational maps and iterative root-finding algorithms*, Ann. of Math. (2) **125** (1987), no. 3, 467–493. MR MR890160 (88i:58082)
- [20] Clayton Petsche, Lucien Szpiro, and Michael Tepper, *Isotriviality is equivalent to potential good reduction for endomorphisms of \mathbb{P}^n over function fields*, Journal of Algebra **322** (2009), 3345–3365.

- [21] Richard Pink, *Finiteness and liftability of postcritically finite quadratic morphisms in arbitrary characteristic*, arXiv:1305.2841, 2013.
- [22] Joseph H. Silverman, *The space of rational maps on \mathbf{P}^1* , Duke Math. J. **94** (1998), no. 1, 41–77. MR MR1635900 (2000m:14010)
- [23] ———, *The arithmetic of dynamical systems*, Graduate Texts in Mathematics, no. 241, Springer-Verlag, New York, 2007.
- [24] Tamás Szamuely, *Galois groups and fundamental groups*, first ed., Cambridge Studies in Advanced Mathematics, vol. 117, Cambridge University Press, Cambridge, 2009.