

Problem Set 3

Sum of Squares Seminar

September 23, 2017, Due October 2, 2017

Problem 1: SOS Proofs (20 points)

Give sum of squares proofs for the following facts (over \mathbb{R}):

- (a) 5 points: $\forall x, y, z, w, 4xyzw \leq x^4 + y^4 + z^4 + w^4$ (this is essentially the AM-GM inequality on 4 terms)
- (b) 5 points: If $\sum_{i=1}^n x_i^2 = 1$ then $\sum_{i=1}^n x_i \leq \sqrt{n}$
- (c) 10 points: If $\sum_{i=1}^n x_i^2 = 1$ then $\prod_{i=1}^n x_i \leq n^{-n}$ (hint is available)
- (d) Challenge question: If M is a doubly stochastic $n \times n$ matrix (i.e. all entries are nonnegative, all rows sum to one, and all columns sum to one), then the permanent of M is at least $\frac{n!}{n^n}$

Problem 2: Decomposing an L^1 pseudo-metric space (15 points)

Recall the objective function for the relaxation of sparsest cut:

$$\frac{\sum_{i < j, (i,j) \in E(G)} d_{ij}}{\sum_{i < j} d_{ij}}$$

Let G be the cycle on 6 vertices, i.e. $V(G) = v_1, \dots, v_6$ and $E(G) = \{(v_i, v_{i+1}) : i \in [1, 5]\} \cup \{(v_1, v_6)\}$. Assume that we are given the following mapping of v_1, \dots, v_6 into \mathbb{R}^2 :

$$v_1 = (0, 1), v_2 = (1, 0), v_3 = (2, 0), v_4 = (3, 0), v_5 = (3, 2), v_6 = (0, 2)$$

- (a) 5 points: What is the value of the objective function given by this L^1 metric? What is the actual sparsity of G ?
- (b) 10 points: Decompose this L^1 metric as a positive linear combination of cut spaces. Which cut space(s) give the best value for the objective function?

Problem 3: Degree 4 Motzkin Polynomial Analogue (15 points)

Consider the polynomial $p(x, y, z) = x^2y^2 + x^2z^2 + y^2z^2 - 4xyz + 1$.

- (a) 5 points: Prove that $\forall x, y, z, p(x, y, z) \geq 0$
- (b) 10 points: Prove that $p(x, y, z)$ cannot be written as the sum of squares of polynomials

Problem 4: SOS Proof of Cauchy-Schwarz with Expected Values (25 points)

Recall the Cauchy-Schwarz inequality: $(\sum_{i=1}^n x_i y_i)^2 \leq (\sum_{i=1}^n x_i^2) (\sum_{i=1}^n y_i^2)$. In lecture, we saw an SOS proof of one analogous statement about pseudo-expectation values, namely

$$\tilde{E} \left[\left(\sum_{i=1}^n x_i y_i \right)^2 \right] \leq \tilde{E} \left[\left(\sum_{i=1}^n x_i^2 \right) \left(\sum_{i=1}^n y_i^2 \right) \right]$$

In this problem, we prove that there is also a sum of squares proof of the following alternative analogue of Cauchy-Schwarz:

$$\tilde{E} \left[\sum_{i=1}^n x_i y_i \right]^2 \leq \tilde{E} \left[\sum_{i=1}^n x_i^2 \right] \tilde{E} \left[\sum_{i=1}^n y_i^2 \right]$$

- (a) 10 points: Prove that for any pseudo-expectation values \tilde{E} , $(\tilde{E}[xy])^2 \leq \tilde{E}[x^2] \tilde{E}[y^2]$
- (b) 5 points: Deduce that for all i, j ,

$$2\tilde{E}[x_i y_i] \tilde{E}[x_j y_j] \leq \tilde{E}[x_i^2] \tilde{E}[y_j^2] + \tilde{E}[x_j^2] \tilde{E}[y_i^2]$$

- (c) 10 points: Use this to prove that for any pseudo-expectation values \tilde{E} ,

$$\tilde{E} \left[\sum_{i=1}^n x_i y_i \right]^2 \leq \tilde{E} \left[\sum_{i=1}^n x_i^2 \right] \tilde{E} \left[\sum_{i=1}^n y_i^2 \right]$$

(hints are available)

Problem 5: Reasoning Using Rational Functions (25 points)

Consider the constraint $(x^2 + 1)y = z^2$. We can immediately see that $y \geq 0$ as $y = \frac{z^2}{x^2+1}$ and both the numerator and the denominator must be non-negative. In this question, we consider whether the sum of squares hierarchy can capture this reasoning.

- (a) 5 points: Give a sum of squares proof that if we add the constraint $y \leq -c$ (equivalently $y = -c - u^2$) for any $c > 0$ then the constraints are infeasible over \mathbb{R} .
- (b) 20 points: Show that there exist degree 4 pseudo-expectation values \tilde{E} with $\tilde{E}[y] < 0$ which respect the constraint that $(x^2 + 1)y = z^2$ (hint is available).

Hints

1c. More generally, show that $\forall k \in [1, n], \frac{1}{\binom{n}{k}} \sum_{i_1 < \dots < i_k} \prod_{j=1}^k x_{i_j}^2 \leq \frac{1}{n^k}$ by showing that if the inequality holds for k_1 and k_2 then it holds for $k_1 + k_2$ whenever $k_1 + k_2 \leq n$

4a. Consider the pseudo-expectation value of a square whose coefficients are functions of $\tilde{E}[x^2]$, $\tilde{E}[y^2]$, and/or $\tilde{E}[xy]$.

4b. Use part a to show that $2\tilde{E}[x_i y_i] \tilde{E}[x_j y_j] \leq 2\sqrt{\tilde{E}[x_i^2] \tilde{E}[y_i^2] \tilde{E}[x_j^2] \tilde{E}[y_j^2]}$

5. One way to find such pseudo-expectation values is to start with an actual expectation over distribution of solutions and then show that you can change the value of $\tilde{E}[y]$ to be negative. For this, it is useful to choose the distribution to make the diagonal entries very large without making the rest of the matrix too large. For example, if your distribution sets $x = B$, $y = z = 0$ with probability $B^{-3.5}$ where B is a large constant, this contributes B^{-5} to $E[x^4]$ and almost nothing to the expected value of any other degree 4 monomial. Thus, we can make the x^4 entry arbitrarily large with negligible effect on the rest of the matrix.

Alternatively, you can write a semidefinite program to find such pseudo-expectation values.