

A FINITE LOOP SPACE NOT RATIONALLY EQUIVALENT TO A COMPACT LIE GROUP

KASPER K. S. ANDERSEN, TILMAN BAUER, JESPER GRODAL, AND ERIK KJÆR PEDERSEN

ABSTRACT. We construct a connected finite loop space of rank 66 and dimension 1254 whose rational cohomology is not isomorphic as a graded vector space to the rational cohomology of any compact Lie group, hence providing a counterexample to a classical conjecture. Aided by machine calculation we verify that our counterexample is minimal, i.e., that any finite loop space of rank less than 66 is in fact rationally equivalent to a compact Lie group, extending the classical known bound of 5.

1. INTRODUCTION

Since the discovery of the Hilton-Roitberg ‘criminal’ [19, 20, 37] in 1968 it has been clear that not every finite loop space is homotopy equivalent to a compact Lie group. The conjecture emerged however, that this should hold rationally, i.e., that any finite loop space should be rationally equivalent to some compact Lie group (see [1], [24, p. 67]), and evidence for this has been accumulated over the years. In this paper we resolve this conjecture in the negative by exhibiting a concrete finite loop space of rank 66 whose rational cohomology does not agree with that of any compact Lie group. To do this, we first use Sullivan’s arithmetic square [40, 41][6, VI.8.1] and the theory of p -compact groups [10, 13, 3] to translate the conjecture into a purely combinatorial statement. We then proceed to show that this statement is ‘generically’ false, with a counterexample appearing in rank 66. On the other hand we verify using a computer that our counterexample is in fact of minimal rank, i.e., that the conjecture is true for any finite loop space of rank less than 66. This extends earlier work of many authors which show the statement to be true when the rank is at most 5 (see [33, 36, 39, 22, 15] and also [2, 27, 29, 28]).

We now explain this in more detail. Recall that a finite connected loop space is a triple (Y, BY, e) where Y is a finite connected CW-complex, BY is a based CW-complex, and $e : Y \rightarrow \Omega BY$ is a homotopy equivalence, where ΩBY denotes the space of based loops in BY . (We usually refer to a loop space just as Y suppressing the rest of the structure.) It is an old theorem of Hopf [21, Satz I] that the rational cohomology of any connected finite loop space is a graded exterior algebra $H^*(Y; \mathbf{Q}) \cong \bigwedge_{\mathbf{Q}}(x_1, \dots, x_r)$, where the generator x_i is in odd dimension $2d_i - 1$. The number r is called the *rank* of Y and the collection of d_i ’s are called the *degrees* (or if doubled the ‘type’) of Y . It is a classical result of Serre [34] that the collection $\{d_1, \dots, d_r\}$ in fact uniquely determines the rational homotopy type of (Y, BY, e) . The p -completion $(Y_{\hat{p}}, BY_{\hat{p}}, e_{\hat{p}})$ is a connected p -compact group, i.e., $H^*(Y_{\hat{p}}; \mathbf{F}_p)$ is finite dimensional and connected and $BY_{\hat{p}}$ is p -complete (in the sense of Sullivan [40, 41] or Bousfield-Kan [6]; see e.g., [13], [10], or [3] for much more on p -compact groups.)

Date: June 20, 2003.

1991 *Mathematics Subject Classification.* Primary: 55P35; Secondary: 55P15, 55R35.

The third named author was partially supported by NSF grant DMS-0104318.

The fourth named author thanks SFB 478 Münster for its hospitality and support.

An amazing result of Dwyer-Wilkerson [13], extending work of Dwyer-Miller-Wilkerson [11] and Adams-Wilkerson [1], says that any p -compact group X has a maximal torus, that is a loop map $T \cong (S^1_p)^r \rightarrow X$ which is suitably maximal and an associated Weyl group W_X . If X is connected, then W_X acts faithfully on $L_X = \pi_1(T)$, in such a way that (W_X, L_X) becomes a finite $\mathbf{Z}_p$ -reflection group and

$$H^*(BX; \mathbf{Z}_p) \otimes \mathbf{Q} \xrightarrow{\cong} (H^*(BT; \mathbf{Z}_p) \otimes \mathbf{Q})^{W_X}.$$

The invariant ring $(H^*(BT; \mathbf{Z}_p) \otimes \mathbf{Q})^{W_X}$ is a polynomial algebra with generators in dimensions $2e_1, \dots, 2e_r$, and the integers $e_1, \dots, e_r$ are just the well known degrees of the $\mathbf{Q}_p$ -reflection group $(W_X, L_X \otimes \mathbf{Q})$ [5, Ch. 7], [17]. (In fact the harder classification states that (W_X, L_X) completely classifies X when p is odd [3].) If Y is a finite loop space then, for all primes p ,

$$H^*(BY; \mathbf{Q}) \otimes \mathbf{Q}_p \cong H^*(BY; \mathbf{Z}) \otimes \mathbf{Q}_p \cong H^*(BY_p^\wedge; \mathbf{Z}_p) \otimes \mathbf{Q}.$$

Furthermore by the Eilenberg-Moore spectral sequence $H^*(BY; \mathbf{Q})$ is a polynomial algebra with generators in dimensions $2d_1, \dots, 2d_r$, where the d_i 's are the degrees of Y introduced earlier. Hence by the above we conclude that, for each prime p , $Y_p^\wedge$ has to be some p -compact group such that the degrees of Y match up with the degrees of the $\mathbf{Q}_p$ -reflection group $(W_{Y_p^\wedge}, L_{Y_p^\wedge} \otimes \mathbf{Q})$. This puts severe restrictions on the possible degrees. Finite $\mathbf{Q}_p$ -reflection groups have been classified by Clark-Ewing [9] building on the classification over $\mathbf{C}$ by Shephard-Todd [35]. The classification divides into three infinite families along with 34 sporadic cases (the non-Lie ones only being realizable for certain primes). We denote by W_D the Weyl group coming from the Dynkin diagram D whereas the notation $G(\cdot, \cdot, \cdot)$ means a given group from the infinite family 2, and G_n refers to one of the other exotic cases, in the standard notation listed e.g., in [9] or [17]. Historically, pioneering work of Clark [8] had already shown that if the maximal degree of Y is h then Y also has to have the degree $m \leq h$ if $m - 1$ and h are relatively prime, using arguments only involving large primes (compare also [23, 3.20]). This, as input to small rank calculations, served as original motivation for the conjecture. Adams-Wilkerson [1] much later found the restrictions imposed by reflection groups described above, but worked only at large primes since the technology of [13] was not available. They furthermore gave an example ([1, Ex. 1.4]) showing that the large prime information is algebraically not enough to settle the conjecture. What we show here is that, contrary to general expectation, the restrictions at all primes are not even sufficient.

Theorem 1.1. *There exists a connected finite loop space Y of rank 66 such that $Y_p^\wedge$ has Weyl group as a $\mathbf{Q}_p$ -reflection group given by*

$$\begin{aligned} W_{A_4 B_4 B_5 B_8 B_8 E_8 A_{12} B_{14}} \times G_{24} & \quad \text{for } p = 2, \\ W_{A_4 D_4 B_5 B_8 B_8 E_8 A_{13} B_{14}} \times G_{12} & \quad \text{for } p \equiv 1, 3 \pmod{8}, \\ W_{G_2 A_4 B_4 B_4 B_7 D_{10} A_{13} B_{15}} \times G(4, 2, 7) & \quad \text{for } p \equiv 5 \pmod{8}, \\ W_{A_4 B_4 B_5 B_8 B_8 E_8 A_{13} B_{14}} \times G(6, 3, 2) & \quad \text{for } p \equiv 7 \pmod{24}, \\ W_{D_4 A_5 D_8 D_8 B_{10} A_{13} D_{16}} \times G(24, 24, 2) & \quad \text{for } p \equiv 23 \pmod{24}. \end{aligned}$$

The space Y has dimension 1254 and the degrees of Y are

$$\{2^8, 3^2, 4^8, 5^2, 6^7, 7, 8^7, 9, 10^5, 11, 12^5, 13, 14^5, 16^3, 18^2, 20^2, 22, 24^2, 26, 28, 30\}$$

using exponent notation to denote repeated degrees, and these do not agree with the degrees of any $\mathbf{Q}$ -reflection group, i.e., the graded vector space $H^*(X; \mathbf{Q})$ does not agree with $H^*(G; \mathbf{Q})$ for any compact Lie group G .

Furthermore this counterexample is minimal in the sense that any connected finite loop space of rank less than 66 is rationally equivalent to some compact Lie group G .

A p -compact group realizing each of the above simple non-Lie rational Weyl groups was constructed by Clark-Ewing [9] in the cases where p does not divide the Weyl group order. The remaining important small prime cases were constructed by Zabrodsky ($G_{12}, p = 3$) [46], Dwyer-Wilkerson ($G_{24}, p = 2$) [12], and Notbohm-Oliver ($G(\cdot, \cdot, \cdot), p$ small) [32]. Since G_{24} and G_{12} are the only finite simple $\mathbf{Q}_p$ -reflection groups which do not come from compact Lie groups, for $p = 2$ and 3 respectively, any counterexample will have to involve these groups.

We note that by work of Bauer-Kitchloo-Notbohm-Pedersen [4] the loop space Y of Theorem 1.1 is in fact homotopy equivalent to a compact, smooth, parallelizable manifold.

Sullivan's arithmetic square [40, 41] reduces the study of finite loop spaces to the study p -compact groups for all primes p with the same degrees together with the well understood concept of arithmetic square 'mixing'. We restrict ourselves here to giving the following lemma which guarantees that an algebraic counterexample produces a topological counterexample.

Lemma 1.2. *Let $\{d_1, \dots, d_r\}$ be a collection of positive integers (with repetitions allowed). Suppose that for each prime p we have a connected p -compact group X_p whose Weyl group (W_{X_p}, L_{X_p}) has degrees $\{d_1, \dots, d_r\}$. Then there exists a (non-unique) connected finite loop space Y such that $\hat{Y}_p \cong X_p$ as p -compact groups.*

The next theorem guarantees that 'generically' there will be sets of degrees which are the degrees of a $\mathbf{Q}_p$ -reflection group for all primes p without being the degrees of any $\mathbf{Q}$ -reflection group. Together with Lemma 1.2 this shows why examples like the one in Theorem 1.1 exist. For the statement of the result we need to introduce some more notation. If $\{d_1, d_2, \dots, d_r\}$ is a collection of degrees, then the associated *degree vector* equals $(x_1, x_2, \dots) \in \mathbf{Z}^{(\infty)}$ where x_i is the number of degrees equal to i . We let $K_{\text{Lie}} \subseteq \mathbf{Q}^{(\infty)}$ denote the positive rational cone spanned by the degree vectors of the simple $\mathbf{Q}$ -reflection groups, i.e. the set of finite nonnegative rational linear combinations of these vectors. Similarly we let K_p denote the positive rational cone spanned by the degree vectors of the simple $\mathbf{Q}_p$ -reflection groups. Finally we let K_{Lin} denote the positive rational cone spanned by the degree vectors of the simple $\mathbf{Q}$ -reflection groups and the degree vectors of the groups $G(m, m, 2)$, $m = 8, 12, 24$, cf. [27, Thm. 1.1(b)].

Theorem 1.3. *We have $K_{\text{Lie}} \subsetneq \bigcap_p K_p$, where the intersection is taken over all primes p . Moreover $\bigcap_p K_p = K_2 \cap K_3 \cap K_5 \cap K_7 \cap K_{\text{Lin}}$.*

Acknowledgments. We are grateful to Ib Madsen whose encouraging questions about integrality results in light of the p -complete theory led us to consider this conjecture. We thank Jørgen Tind for pointing us to Komei Fukuda's program `cdd+` [16] for doing computations on polyhedral cones. The hospitality of University of Copenhagen, SFB 478 Münster, and Århus University helped make this collaboration possible.

2. PROOFS

Recall the following essentially classical result.

Theorem 2.1. *Let X be a simply connected p -compact group with degrees $\{d_1, \dots, d_r\}$. Then, as a space, $X \simeq (S^{2d_1-1} \times \dots \times S^{2d_r-1})_p^\wedge$ if and only if $p \geq \max\{d_1, \dots, d_r\}$. If X is just assumed connected then under the stronger assumption $p > \max\{d_1, \dots, d_r\}$, X still splits as a product of spheres.*

Sketch of proof. Set $h = \max\{d_1, \dots, d_r\}$ and suppose first that X is simply connected. If $p \geq h$ then by a Bockstein spectral sequence argument of Browder [7, Thm. 4.7], $H^*(X; \mathbf{Z}_p)$ is torsion free and concentrated in odd degrees. Hence an easy argument of Serre [34, Ch. V Prop. 6] (see also [26]), using that $\pi_n(S^{2d_i-1})$ has no p -torsion when $n < 2d_i - 1 + 2p - 3$, yields that $X \simeq (S^{2d_1-1} \times \dots \times S^{2d_r-1})_p^\wedge$. (In fact, this direction uses only that X is an H -space with $H^*(X; \mathbf{F}_p)$ finite dimensional.) The other direction, which is more subtle and not needed here, was first established for compact Lie groups by Serre [34] and Kumpel [25] by case-by-case arguments, and later a general argument was given by Wilkerson [44, Thm. 4.1] using operations in K -theory.

Assume now that X is just connected and that $p > h$. If $\pi_1 X$ is torsion free, then as a space $X \simeq \tilde{X} \times (S^1_p)^\wedge$ where $\tilde{X}$ is simply connected (cf. e.g. [24, p. 24]), which reduces us to the previous case. Hence we just have to justify that with p as above $\pi_1(X)$ does not have torsion. By [30, Thm. 1.4] we have a fibration $BK \rightarrow B\tilde{X} \times BT' \rightarrow BX$ such that $\tilde{X}$ is a simply connected p -compact group, T' is a torus, K is a finite p -group, and the projection map $K \rightarrow \tilde{X}$ is a central monomorphism. But then by [14, Thm. 7.6] K is contained in $\check{T}^{W_{\tilde{X}}}$, where $\check{T}$ is a discrete approximation to maximal torus in $\tilde{X}$. In particular if $\check{T}^{W_{\tilde{X}}} = 0$, $\pi_1(X)$ has to be torsion free. But if $p > h$ then in particular $p \nmid |W_{\tilde{X}}|$, so we have an exact sequence

$$\dots \rightarrow H^0(W_{\tilde{X}}; L_{\tilde{X}} \otimes \mathbf{Q}) \rightarrow H^0(W_{\tilde{X}}; \check{T}) \rightarrow H^1(W_{\tilde{X}}; L_{\tilde{X}}) \rightarrow \dots$$

where the first and third terms are zero so $\check{T}^{W_{\tilde{X}}} = 0$ as wanted. $\square$

Proof of Lemma 1.2. Set $h = \max\{d_1, \dots, d_r\}$ and let $BM = (\prod_p BX_p)_{\mathbf{Q}}$. Since rationalization commutes with taking loop space and finite products we have that $\Omega BM \simeq (\prod_{p < h} (X_p)_{\mathbf{Q}}) \times (\prod_{p \geq h} X_p)_{\mathbf{Q}}$. By Theorem 2.1 $X_p \simeq (S^{2d_1-1} \times \dots \times S^{2d_r-1})_p^\wedge$ when $p > h$ and by the same argument $(X_p)_{\mathbf{Q}} \simeq ((S^{2d_1-1} \times \dots \times S^{2d_r-1})_p^\wedge)_{\mathbf{Q}}$ for all primes p . Combined with the fact that $(S^{2d_i-1}_p^\wedge)_{\mathbf{Q}} \simeq K(\mathbf{Q}_p, 2d_i - 1)$ this implies that $\pi_n(BM) = (\prod_n \pi_n(BX_p))_{\mathbf{Q}} \cong \bigoplus_{i, 2d_i=n} \mathbf{A}_f$, where $\mathbf{A}_f = (\prod_p \mathbf{Z}_p) \otimes \mathbf{Q}$ is the ring of finite adeles. In particular BM only has homotopy groups in even dimensions. But now a rational space which only has homotopy groups in even dimensions is necessarily a product of Eilenberg-Mac Lane spaces, as is easily seen by going up the Postnikov tower (cf. e.g., [43, Ch. IX]). Set $BK = K(\mathbf{Q}, 2d_1) \times \dots \times K(\mathbf{Q}, 2d_r)$ and construct a map $BK \rightarrow BM$ by levelwise taking the unit ring map $\mathbf{Q} \rightarrow \mathbf{A}_f$. Define BY as the homotopy pullback of the diagram $BK \rightarrow BM \leftarrow \prod_p BX_p$.

Since $\mathbf{Q}$ and $\hat{\mathbf{Z}} = \prod_p \mathbf{Z}_p$ generate $\mathbf{A}_f$ the Mayer-Vietoris sequence in homotopy groups corresponding to a homotopy pull-back in fact splits hence $\pi_n(BY)$ is the pull-back in groups of the diagram $\pi_n(BK) \rightarrow \pi_n(BM) \leftarrow \prod_p \pi_n(BX_p)$. Concretely, the homotopy groups of BY are given by

$$\pi_n(BY) = \left(\bigoplus_{i, 2d_i=n} \mathbf{Z} \right) \oplus \left(\bigoplus_p \text{Tor}(\mathbf{Z}, \pi_n(BX_p)) \right).$$

In particular, this shows that $\pi_n(\Omega BY)$ is finitely generated for all n . Hence also $H^n(\Omega BY; \mathbf{Z})$ is finitely generated for all n (see [34][18, Thm. 2.16]).

By construction $H^*(BY; \mathbf{Z}_p) \cong H^*(BX_p; \mathbf{Z}_p)$ so, since the spaces involved are simply connected, $H^*(\Omega BY; \mathbf{Z}_p) \cong H^*(X_p; \mathbf{Z}_p)$ for all p . But since we have seen that each $H^n(\Omega BY; \mathbf{Z}_p)$ is finitely generated and we know that X_p is homotopy equivalent to a product of spheres for all $p > h$, we conclude that in fact $\bigoplus_n H^n(\Omega BY; \mathbf{Z})$ is finitely generated as an abelian group.

If ΩBY is simply connected then it follows from the classical results of Wall [42, Thm. B+F] that ΩBY is weakly homotopy equivalent to a finite CW-complex Y . If ΩBY is not simply connected then the conclusion still holds, now appealing to a more recent result of Notbohm [31] (see also [4]) which relies on ΩBY being a loop space. $\square$

Proof of Theorem 1.1. It follows directly from Lemma 1.2 that we can construct a connected finite loop space Y with the listed properties. One can check directly by a finite search that the degrees of Y does not agree with those of a compact Lie group, but one can also argue more simply as follows. There are exactly 61 simple $\mathbf{Q}$ -reflection groups whose degrees are all at most 30. The inner product of the vector

$$(2.1) \quad (0, 2, -1, -1, 0, 0, 1, -1, 0, 0, 0, 1, -1, 0, 1, -1, 0, 0, 0, 0, 0, 4, -1, -3, 1, -1, 0, 0, 0, 1)$$

and the degree vector of any of these is non-negative. Hence the same holds for any $\mathbf{Q}$ -reflection groups whose degrees are all at most 30. However the inner product of the vector (2.1) with the degree vector of Y equals -1 , so the degrees of Y does not agree with those of a compact Lie group.

To check that our counterexample has minimal rank we proceed as follows. For any prime p , there are only finitely many $\mathbf{Q}_p$ -reflection groups of a given rank, cf. [9]. Hence one can go through the list of (say) $\mathbf{Q}_2$ -reflection groups and check which of these has degrees not matching those of any $\mathbf{Q}$ -reflection group, but matching those of a $\mathbf{Q}_p$ -reflection group for $p = 3, 5, \dots$. We have written a C++ program which implements this algorithm, and used it to check all $\mathbf{Q}_2$ -reflection groups of rank less than 66. $\square$

Remark 2.2. Note that Theorem 1.1 in particular tells us that there exists a loop space whose p -completion is not homotopy equivalent to the p -completion of a compact Lie group for any prime p . If we only want this to hold for a single prime p we can find much simpler examples. For instance using Lemma 1.2 one can construct a finite loop space which 2-completed is homotopy equivalent to $X_2 = DI(4) \times \mathrm{Sp}(1)_2 \times \mathrm{Sp}(6)_2$ and which p -completed for $p \neq 2$ is homotopy equivalent to $X_p = \mathrm{Sp}(3)_p \times \mathrm{Sp}(7)_p$. However, X_2 is not homotopy equivalent to the 2-completion of a compact Lie group, since the only Lie groups with the right rational degrees are quotients of $\mathrm{Sp}(3) \times \mathrm{Sp}(7)$, but these do not have the same mod 2 Poincaré series as X_2 , as can be obtained from [12]. (Compare [45, Conj. 2], [38, Prob. 9], [24, Conj. B+C].)

Proof of Theorem 1.3. The first claim follows from the proof of Theorem 1.1, see also Remark 2.3 below. To show the second claim note that the proof of [27, Thm. 1.1(b)] shows that $\bigcap_p K_p \subseteq K_{\mathrm{Lin}}$ and hence

$$\bigcap_p K_p \subseteq K_2 \cap K_3 \cap K_5 \cap K_7 \cap K_{\mathrm{Lin}}.$$

To prove the reverse inclusion, note that the only simple $\mathbf{Q}_3$ -reflection group which is not a $\mathbf{Q}$ -reflection group is the group G_{12} . Since this is a $\mathbf{Q}_p$ -reflection group for all primes p satisfying $p \equiv 1, 3 \pmod{8}$ we get $K_3 \subseteq K_p$ for these primes. Similarly the simple $\mathbf{Q}_5$ -reflection groups which are not $\mathbf{Q}$ -reflection groups are precisely the groups $G(4, 1, n), G(4, 2, n)$ for

$n \geq 2$, $G(4, 4, n)$ for $n \geq 3$, $\mathbf{Z}/4$ from family 3, G_8 , G_{29} and G_{31} . This shows that $K_5 \subseteq K_p$ when $p \equiv 1 \pmod{4}$. In the same way we see that $K_7 \subseteq K_p$ when p satisfies $p \equiv 1 \pmod{6}$ and $p \equiv \pm 1 \pmod{8}$, i.e. when $p \equiv 1, 7 \pmod{24}$. Finally the groups $G(m, m, 2)$, $m = 8, 12, 24$ are all $\mathbf{Q}_p$ -reflection groups when $p \equiv \pm 1 \pmod{24}$, so $K_{\text{Lin}} \subseteq K_p$ for these primes. This proves the result since any prime p satisfies $p = 2$, $p \equiv 1, 3 \pmod{8}$, $p \equiv 1 \pmod{4}$, $p \equiv 1, 7 \pmod{24}$ or $p \equiv \pm 1 \pmod{24}$. $\square$

Remark 2.3. A few remarks about how we found the counterexample in Theorem 1.1 might be in order, since this is not really clear from the proof. First we used Fukuda's cdd+ program [16] to establish Theorem 1.3 by showing that K_{Lie} truncated at degree say 30 does not agree with the intersections of the similarly truncated versions of K_2 , K_3 , K_5 , K_7 , and K_{Lin} . From this it is a linear programming problem to obtain a concrete point in the difference, and by solving the associated integer programming problem one gets a point in the difference which is minimal with respect to rank (or dimension). Note however that being minimal in this sense is slightly weaker than being a minimal counterexample, which is why we had to finish off our proof of minimality in Theorem 1.1 with a brute force check, which required rather massive computer calculations. Using the geometric picture we have found a counterexample of smaller dimension but larger rank. Namely, there exist a connected finite loop space with rank 74 and dimension 1250 and degrees

$$\{2^9, 3^2, 4^7, 5^3, 6^8, 7^3, 8^8, 9^3, 10^6, 11^2, 12^6, 13^2, 14^5, 15, 16^3, 18^2, 20^2, 22, 24\},$$

which is not rationally equivalent to a compact Lie group. This is seen by considering the $\mathbf{Q}_p$ -reflection groups

$$\begin{aligned} W_{A_1 A_1 E_6 D_7 D_8 A_9 D_{11} D_{13} A_{15}} &\times G_{24} && \text{for } p = 2, \\ W_{A_1 A_1 D_5 D_7 D_9 B_{10} B_{12} A_{13} A_{14}} &\times G_{12} && \text{for } p \equiv 1, 3 \pmod{8}, \\ W_{A_1 A_1 D_5 D_6 E_7 D_9 B_{11} A_{13} A_{14}} &\times G(4, 4, 7) && \text{for } p \equiv 5 \pmod{8}, \\ W_{G_2 D_5 D_7 D_9 B_{10} B_{12} A_{13} A_{14}} &\times G(8, 8, 2) && \text{for } p \equiv 7 \pmod{8}. \end{aligned}$$

It is also possible to construct a counterexample where all degrees are even, cf. [29]. For instance there is an example of rank 68, dimension 1468 and degrees

$$\{2^8, 4^8, 6^8, 8^8, 10^6, 12^5, 14^6, 16^4, 18^3, 20^3, 22^2, 24^3, 26^2, 28, 30\}.$$

Here one can use the $\mathbf{Q}_p$ -reflection groups

$$\begin{aligned} W_{B_4 B_5 B_5 B_8 B_8 E_8 B_{13} B_{14}} &\times G_{24} && \text{for } p = 2, \\ W_{D_4 B_5 B_5 B_8 B_8 E_8 B_{14} D_{14}} &\times G_{12} && \text{for } p \equiv 1, 3 \pmod{8}, \\ W_{G_2 B_4 B_5 B_5 B_7 B_9 D_{14} B_{15}} &\times G(4, 2, 7) && \text{for } p \equiv 5 \pmod{8}, \\ W_{B_4 B_5 B_5 B_8 B_8 E_8 B_{14} D_{14}} &\times G(6, 3, 2) && \text{for } p \equiv 7 \pmod{24}, \\ W_{D_4 D_6 D_8 D_8 B_{10} D_{14} D_{16}} &\times G(24, 24, 2) && \text{for } p \equiv 23 \pmod{24}. \end{aligned}$$

REFERENCES

- [1] J. F. Adams and C. W. Wilkerson. Finite H -spaces and algebras over the Steenrod algebra. *Ann. of Math. (2)*, 111(1):95–143, 1980. (Erratum: *Ann. of Math. (2)* **113** (1981), no. 3, 621–622).
- [2] J. Aguadé. A note on realizing polynomial algebras. *Israel J. Math.*, 38(1-2):95–99, 1981.
- [3] Kasper K. S. Andersen, Jesper Grodal, Jesper M. Møller, and Antonio Viruel. The classification of p -compact groups for p odd. arXiv:math.AT/0302346.
- [4] Tilman Bauer, Nitya Kitchloo, Dietrich Notbohm, and Erik K. Pedersen. Finite loop spaces are manifolds. Preprint; available from <http://www.math.binghamton.edu/erik>, 2003.

- [5] D. J. Benson. *Polynomial invariants of finite groups*. Cambridge University Press, Cambridge, 1993.
- [6] A. K. Bousfield and D. M. Kan. *Homotopy limits, completions and localizations*. Springer-Verlag, Berlin, 1972. Lecture Notes in Mathematics, Vol. 304.
- [7] William Browder. On differential Hopf algebras. *Trans. Amer. Math. Soc.*, 107:153–176, 1963.
- [8] Allan Clark. On π_3 of finite dimensional H -spaces. *Ann. of Math. (2)*, 78:193–196, 1963.
- [9] Allan Clark and John Ewing. The realization of polynomial algebras as cohomology rings. *Pacific J. Math.*, 50:425–434, 1974.
- [10] W. G. Dwyer. Lie groups and p -compact groups. In *Proceedings of the International Congress of Mathematicians, Extra Vol. II (Berlin, 1998)*, pages 433–442 (electronic), 1998.
- [11] W. G. Dwyer, H. R. Miller, and C. W. Wilkerson. Homotopical uniqueness of classifying spaces. *Topology*, 31(1):29–45, 1992.
- [12] W. G. Dwyer and C. W. Wilkerson. A new finite loop space at the prime two. *J. Amer. Math. Soc.*, 6(1):37–64, 1993.
- [13] W. G. Dwyer and C. W. Wilkerson. Homotopy fixed-point methods for Lie groups and finite loop spaces. *Ann. of Math. (2)*, 139(2):395–442, 1994.
- [14] W. G. Dwyer and C. W. Wilkerson. The center of a p -compact group. In *The Čech centennial (Boston, MA, 1993)*, pages 119–157. Amer. Math. Soc., Providence, RI, 1995.
- [15] John Ewing. On the type of associative H -spaces. *Bull. Amer. Math. Soc.*, 78:35–37, 1972.
- [16] Komei Fukuda. *cdd and cddplus Homepage*. McGill University, Montreal, Canada, April 2003. http://www.cs.mcgill.ca/~fukuda/soft/cdd_home/cdd.html.
- [17] M. Geck and G. Malle. Reflection groups. To appear in *Handbook of Algebra*, vol. 4, North-Holland, Amsterdam, 2003.
- [18] Peter Hilton, Guido Mislin, and Joe Roitberg. *Localization of nilpotent groups and spaces*. North-Holland Publishing Co., Amsterdam, 1975. North-Holland Mathematics Studies, No. 15, Notas de Matemática, No. 55. [Notes on Mathematics, No. 55].
- [19] Peter Hilton and Joseph Roitberg. Note on principal S^3 -bundles. *Bull. Amer. Math. Soc.*, 74:957–959, 1968.
- [20] Peter Hilton and Joseph Roitberg. On principal S^3 -bundles over spheres. *Ann. of Math. (2)*, 90:91–107, 1969.
- [21] Heinz Hopf. Über die Topologie der Gruppen-Mannigfaltigkeiten und ihre Verallgemeinerungen. *Ann. of Math. (2)*, 42:22–52, 1941.
- [22] J. R. Hubbuck. Associative H -spaces with small ranks. *Proc. Amer. Math. Soc.*, 30:375–382, 1971.
- [23] James E. Humphreys. *Reflection groups and Coxeter groups*. Cambridge University Press, Cambridge, 1990.
- [24] Richard M. Kane. *The homology of Hopf spaces*. North-Holland Publishing Co., Amsterdam, 1988.
- [25] P. G. Kumpel, Jr. Lie groups and products of spheres. *Proc. Amer. Math. Soc.*, 16:1350–1356, 1965.
- [26] P. G. Kumpel, Jr. On p -equivalences of mod p H -spaces. *Quart. J. Math. Oxford Ser. (2)*, 23:173–178, 1972.
- [27] James P. Lin. The rational cohomology of finite loop spaces. *J. Pure Appl. Algebra*, 53(1-2):71–73, 1988.
- [28] James P. Lin. Cup products and finite loop spaces. *Topology Appl.*, 45(1):73–84, 1992.
- [29] James P. Lin and Frank Williams. The type of a torsion free finite loop space. *Topology Appl.*, 42(2):175–186, 1991.
- [30] J. M. Møller and D. Notbohm. Centers and finite coverings of finite loop spaces. *J. Reine Angew. Math.*, 456:99–133, 1994.
- [31] D. Notbohm. The finiteness obstruction for loop spaces. *Comment. Math. Helv.*, 74(4):657–670, 1999.
- [32] Dietrich Notbohm. Topological realization of a family of pseudoreflection groups. *Fund. Math.*, 155(1):1–31, 1998.
- [33] Shōji Ochiai. On the type of an associative H -space of rank three. *Proc. Japan Acad.*, 44:811–815, 1968.
- [34] Jean-Pierre Serre. Groupes d’homotopie et classes de groupes abéliens. *Ann. of Math. (2)*, 58:258–294, 1953.
- [35] G. C. Shephard and J. A. Todd. Finite unitary reflection groups. *Canadian J. Math.*, 6:274–304, 1954.
- [36] Larry Smith. On the type of an associative H -space of rank two. *Tôhoku Math. J. (2)*, 20:511–515, 1968.
- [37] James D. Stasheff. Manifolds of the homotopy type of (non-Lie) groups. *Bull. Amer. Math. Soc.*, 75:998–1000, 1969.

- [38] James D. Stasheff. H -space problems. In *H-spaces (Actes Réunion, Neuchâtel, 1970)*, pages 122–136. Lecture Notes in Mathematics, Vol. 196. Springer, Berlin, 1971.
- [39] Tamio Sugawara. On the type of an associative H -spaces of rank four. *Mem. Fac. Sci. Kyushu Univ. Ser. A*, 25:191–196, 1971.
- [40] Dennis Sullivan. Geometric topology, Part I, Localization, periodicity and Galois symmetry. Mimeographed notes, MIT, 1970.
- [41] Dennis Sullivan. Genetics of homotopy theory and the Adams conjecture. *Ann. of Math. (2)*, 100:1–79, 1974.
- [42] C. T. C. Wall. Finiteness conditions for CW-complexes. *Ann. of Math. (2)*, 81:56–69, 1965.
- [43] George W. Whitehead. *Elements of homotopy theory*, volume 61 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1978.
- [44] Clarence Wilkerson. K -theory operations in mod p loop spaces. *Math. Z.*, 132:29–44, 1973.
- [45] Clarence Wilkerson. Rational maximal tori. *J. Pure Appl. Algebra*, 4:261–272, 1974.
- [46] A. Zabrodsky. On the realization of invariant subgroups of $\pi_*(X)$. *Trans. Amer. Math. Soc.*, 285(2):467–496, 1984.

DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF COPENHAGEN, UNIVERSITETSPARKEN 5
 DK-2100 COPENHAGEN Ø, DENMARK
E-mail address: `kksa@math.ku.dk`

SONDERFORSCHUNGSBEREICH 478 GEOMETRISCHE STRUKTUREN IN DER MATHEMATIK
 WESTFÄLISCHE WILHELMS-UNIVERSITÄT MÜNSTER, HITTORFSTR. 27
 D-48149 MÜNSTER, GERMANY
E-mail address: `tbauer@math.uni-muenster.de`

DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF CHICAGO, 5734 S. UNIVERSITY AVE.
 CHICAGO, IL 60637, USA
E-mail address: `jg@math.uchicago.edu`

DEPT. OF MATHEMATICAL SCIENCES
 BINGHAMTON UNIVERSITY
 BINGHAMTON, NEW YORK 13902-6000, USA
E-mail address: `erik@math.binghamton.edu`